

**MOOREBANK PRECINCT EAST STAGE 2:
BIODIVERSITY MONITORING IN ANZAC CREEK**

SPRING 2025 SURVEY



**Final Report Prepared for
ARCADIS**

17 February 2026



BIO-ANALYSIS Pty Ltd

Marine & Freshwater Ecology

2/1 Botham Close, Charmhaven, NSW 2263

(Mobile) 0414 477 066; (Email) info@bioanalysis.com.au

Document Information

Project Name	Moorebank Precinct 2 East Stage 2: Biodiversity Monitoring in Anzac Creek (Spring 2025 Survey)
Prepared for	Arcadis Australia Pacific
Prepared by	Dr Sharon Cummins
Reviewed by	Dr Dan Roberts
File Name:	MPES2 B106 Anzac Creek Monitoring Report Spring 2025
Citation	BIO-ANALYSIS (2026). MPES2 B106 Anzac Creek Monitoring Report- Spring 2025.
Cover Photo	Anzac Creek @ Site AQ1, 1 December 2025

Contact Information

Dr Sharon Cummins
Telephone: 043 811 2962
Email: cumminssharon@bigpond.com

Dr Dan Roberts
Telephone: 0414 477 066
Email: dan@bioanalysis.com.au

Document Control

Version	Date Issued
Draft Version 01	7/02/2026
Revised Draft	16/02/2026
Final Version	17/02/2026

Acknowledgements

Shane Murray and William Roberts are thanked for work in the field and laboratory. Dan Roberts reviewed the draft report. Jan Roberts, William Roberts, Nick Roberts (BIO-ANALYSIS), Heather Tilley (Arcadis) and Mark Howley (Tactical) provided management support.

EXECUTIVE SUMMARY

Introduction

The Sydney Intermodal Terminal Alliance (SIMTA) received approval for the construction and operation of Stage 2 (the Project) of the Moorebank Precinct East (MPE) Project, which comprises the second stage of development under the MPE Concept Approval (MP10_0193) and approved under Development Approval SSD 7628.

SIMTA was the original applicant for Stage 1 (SSD 6766) and Stage 2 (SSD 7628), under the MPE Concept Approval. The applicant for the SSD 7628 has been updated to “The Trust Company Limited” (ACN 004 027 749). In 2022, LOGOS Property took over the management of the warehouse and distribution facilities, as well as the overall management of the Moorebank Intermodal East and West Precincts. In July 2024, ESR Group acquired the remaining interest in LOGOS, and overall management of the MIP East Precinct, is now the responsibility of ESR Australia & NZ (ESR). Qube Logistics will continue to maintain responsibility for the IMEX and the Rail Link.

The MPE site, including the Project site, is located approximately 27 km south-west of the Sydney Central Business District (CBD) and approximately 26 km west of Port Botany and includes the former Defence National Storage and Distribution Centre (DNSDC) site. The MPE site is situated within the Liverpool Local Government Area (LGA), in Sydney’s Southwest subregion, approximately 2.5 km from the Liverpool City Centre.

The MPE Project involves the development of an intermodal facility including warehouse and distribution facilities, freight village (ancillary site and operational services), stormwater infrastructure, landscaping, servicing and associated works on the eastern side of Moorebank Avenue. Stage 2 of the MPE Project (MPES2) involves the construction and operation of warehousing and distribution facilities on the MPE site and upgrades to approximately 2.1 km of Moorebank Avenue.

Water during construction will be managed in accordance with the currently approved Construction Environmental Management Plan (CEMP) and will be discharged into the sediment (SED) Basins and into Anzac Creek (via DP5 and DP7).

It was also considered likely that runoff from some areas of the MPES2 site would be collected by a vegetated dam situated within Commonwealth Department of Defence land. Flow from this dam enters Anzac Creek upstream of Site AQ14 via a culvert.

A Baseline Aquatic Ecological Monitoring Program (BAEMP) was developed by Biosis Pty Ltd for Arcadis in March 2018, to address CoC B106. The purpose of the BAEMP was to establish baseline stream health and water quality conditions within selected sites along Anzac Creek prior to commencement of Early Works. This was undertaken in autumn 2018. Construction activities commenced soon after.

The baseline monitoring forms the basis for the ongoing Biodiversity Monitoring Strategy (BMS) to assess stream health in accordance with CoC B106, to determine any change in stream health or water quality throughout the life of the Project and to ascertain whether these changes can be attributed to the Project works. The BMS outlines monitoring requirements and includes the Stormwater Monitoring Strategy required by CoC B43 and B44.

BIO-ANALYSIS Pty Ltd was commissioned by Arcadis on behalf of Tactical Group to assess stream health and water quality at six monitoring sites along Anzac Creek (the Study Area) in spring 2025, in accordance with the BMS.

Methods

The BMS focusses on four main indicators: i) aquatic habitat, including riparian habitat, aquatic macrophytes and fish habitat; ii) surface water quality and sediment characteristics; iii) aquatic macroinvertebrates sampled using the Australian River Assessment System (AUSRIVAS) protocol; and iv) fish sampled using a backpack electro-fisher.

The primary aim of monitoring is to determine whether any change in stream health or water quality occur throughout the life of the MPE Project in accordance with the BMS and to ascertain whether these changes can be attributed to the Project works. Should an indicator variable deteriorate below the range for its baseline value, a stream health investigation protocol is to be initiated under the BAEMPs Adaptive Management Plan.

The sampling design included six sites (approximately 100 m in length). Site AQ1 is situated upstream of the MPE Project. Sites AQ4, AQ8, AQ12, AQ13 and AQ14 are situated at increasing distances downstream of the MPE Project. Stream health monitoring is to be done on two occasions within each of autumn and spring.

The results of the spring 2025 monitoring event were compared with those obtained in autumn 2018 (baseline), spring 2018, autumn and spring 2019, autumn and spring 2020, autumn and spring 2021, autumn and spring 2022, autumn and spring 2023, autumn and spring 2024 and autumn 2025 (during construction).

Results

This report presents the results of i) spring 2025 surveys 1 and 2 and ii) comparisons of the findings of the current survey with the Baseline survey (autumn 2018) and subsequent surveys done each autumn and spring.

Within the current reporting period (spring 2025), an unplanned discharge of sediment laden water from the MPE site to Anzac Creek, between Site AQ1 and AQ4, occurred during November 2025. At the time of the second spring survey (1 December 2025), the water was clear and free of sediment.

Riparian and terrestrial vegetation removed along the upstream reaches of Site AQ4 in May/June 2025, associated with the Moorebank Avenue Realignment¹, showed some signs of regeneration. Extensive cover by vegetation within the riparian zone and stream channel upstream and downstream of Site AQ4 continues to contribute stability to the majority of Anzac Creek, including the large refuge pool situated at Site AQ12.

Concentrations of lead in sediments collected at Site AQ1 (range = 100 to 110 mg/kg during spring 2025) continue to exceed the guideline value (50 mg/kg), including at the time of the baseline (91 mg/kg) survey. Copper, nickel and zinc have occasionally exceeded guideline values, but total petroleum hydrocarbons and poly-fluoroalkyl substances (e.g. PFAS and

¹ The riparian vegetation along Anzac Creek at the location of disturbance will be rehabilitated once the Moorebank Avenue Realignment works are completed in 2026.

PFOS), continue to comply. Site AQ1 is situated upstream of potential inputs from the Project, so no additional testing at this site is considered necessary.

Reduced dissolved oxygen levels, elevated nitrogen, aluminium and copper measured at the refuge pool (Site AQ12), including prior to commencement of the Project, have consistently suggested that aquatic habitat and biota within Anzac Creek are influenced by various types of anthropogenic disturbance. Importantly, the data collected to date indicate that there has been no further degradation of water quality since the Project related construction work began.

Over the course of the monitoring programme, the diversity of aquatic macroinvertebrates, Australian River Assessment System (AUSRIVAS) and Stream Invertebrate Grade Number Average Level (SIGNAL2) scores have been relatively low, indicating that the aquatic macroinvertebrate fauna has experienced one or more forms of human impact. Despite this, some pollution tolerant taxa have commonly been identified, including dragonfly, mayfly and caddis fly families. Importantly, comparison of the AUSRIVAS and SIGNAL2 scores between the baseline and construction phase continue to indicate an overall stability in aquatic health.

Altogether, ten species of fish have been collected from within the refuge pool: three native species of gudgeon, two native species of eel, one native galaxiid species, one native cat-fish species and three introduced species (Gambusia, Goldfish and Oriental weatherloach), confirming that the creek does provide some habitat for native species of fish. All of the species caught are common within NSW. No threatened species of fish listed under the *NSW Fisheries Management Act, 1994* or the *Environment Protection and Biodiversity Conservation Act, 1999* have been recorded.

Notably, a dead Longfin eel was observed at Site AQ14 at the time of Spring 2025 Survey 2. A deep lesion was apparent, similar to those caused by Epizootic Ulcerative Syndrome (EUS) (Red Spot Disease).

Conclusions

Examination of the results from the spring 2025 monitoring event found no evidence of changes in the indicator variables (surface water and sediment quality, assemblages of aquatic macroinvertebrates and fish) that could be attributed to the Project works. Thus, in accordance with the Biodiversity Monitoring Strategy, no adaptive management contingency measure was triggered.

Recommendations

It is recommended that the stream health monitoring programme is repeated in autumn and spring 2026 using the methods employed for baseline and operation phase surveys. It is also recommended that Land Managers focus on:

- stabilising areas of the stream bank recently cleared within the vicinity of Site AQ4, to minimise sediments and contaminants becoming mobilised to downstream environments;
- containment and on-going suppression of Alligator Weed and Ludwigia at Site AQ1, and the popular aquarium plant, *Egeria densa* (Egeria), within the refuge pool at Site AQ12. Signage and public information at popular points of entry by the public to the creek and other local waterways may reduce the chance of unintentional human-assisted introductions (e.g. by using live bait, or by being released by aquaria) of aquatic plants and fish.

TABLE OF CONTENTS

EXECUTIVE SUMMARY.....	3
1.0 INTRODUCTION.....	10
2.0 METHODS.....	11
2.1 Study Area.....	11
2.3 Performance Measures and Indicators	14
2.4 Field Methods.....	16
2.4.1 Visual Stream Assessments.....	17
2.4.2 Surface Water Quality & Sediment Monitoring	17
2.4.3 Aquatic Macroinvertebrates	18
2.4.5 Data Analysis	19
2.4.6 Quality Assurance/Quality Control (QA/QC).....	20
3.0 RESULTS.....	21
3.1 Aquatic Habitat Characteristics.....	21
3.2 Water & Sediment Characteristics	28
3.2.1 Water Quality	28
3.2.2 Sediment Characteristics	42
3.3 Aquatic Macroinvertebrates	59
3.4 Fish.....	63
5.0 DISCUSSION	66
5.1 Aquatic Habitat and Hydrology	66
5.2 Water Quality	67
5.3 Biological Monitoring.....	67
Summary	68
6.0 CONCLUSION & RECOMMENDATIONS	69
7.0 REFERENCES.....	70
APPENDICES.....	73
Appendix 1 - GPS positions (UTMs) for stream monitoring sites (spring 2025).....	73
Appendix 2 – Visual Assessment Scores	74
Appendix 3 - Macroinvertebrate taxa collected at Site AQ12 in spring 2025 using the NSW AUSRIVAS protocol. Mosquito fish were also collected in the net samples.....	77

List of Tables

Table 1. Date and information on aquatic ecology monitoring completed for the Project.	Error! Bookmark not defined.
Table 2. Assessment types recommended for each monitoring site (Biosis, 2018).....	15
Table 3. Indicator variables and adaptive management contingency measures.....	16
Table 4. Mean (+ SE) physico-chemical water quality and nutrient values recorded at the time of the Baseline (autumn 2018, $n = 1$) and the autumn 2024 ($n = 3$) surveys and the appropriate Default Trigger Values (DTV). Values highlighted in bold type indicate where results were outside the recommended DTV.....	30
Table 6. Summary of dissolved metal compound results for Site AQ12 between autumn 2018 (Baseline) and autumn 2024 ($n = 1$).....	31
Table 7. Summary of BTEX and perfluorinated compound results ($n = 1$). Error! Bookmark not defined.	
Table 8. Mean (\pm SE) sediment metal results (mg/L) for surveys done between autumn 2018 ($n = 1$) and autumn 2025 ($n = 2$).	
Table 9. Total number of taxa, AUSRIVAS & SIGNAL 2 outputs for Site AQ12 ($n = 1$)	59
Table 10. Fish collected at Site AQ12 between autumn 2018 and autumn 2025.....	64

List of Figures

Figure 1. Project Location.....	12
Figure 2. Rainfall (mm) measured at Holsworthy Aerodrome AWS Rainfall Station (66161) between 1 January and 31 December 2025.....	22
Figure 3. OE50 Taxa Scores and their respective Band Scores (B-D) for AUSRIVAS samples collected at Site AQ12 since autumn 2018. NB Note that the bands displayed are relevant to autumn edge habitat, these being slightly different to spring.	53
Figure 4. Mean (\pm) SIGNAL 2 results for Site AQ12 sampled in Anzac Creek since autumn 2018.....	54

1.0 INTRODUCTION

The Sydney Intermodal Terminal Alliance (SIMTA) received approval for the construction and operation of Stage 2 (the Project) of the Moorebank Precinct East (MPE) Project, which comprises the second stage of development under the MPE Concept Approval (MP10_0193) and approved under Development Approval SSD 7628.


SIMTA was the original applicant for Stage 1 (SSD 6766) and Stage 2 (SSD 7628), under the MPE Concept Approval. The applicant for the SSD 7628 has been updated to “The Trust Company Limited” (ACN 004 027 749). In 2022, LOGOS Property took over the management of the warehouse and distribution facilities, as well as the overall management of the Moorebank Intermodal East and West Precincts. In July 2024, ESR Group acquired the remaining interest in LOGOS, and overall management of the MIP East Precinct, is now the responsibility of ESR Australia & NZ (ESR). Qube Logistics will continue to maintain responsibility for the IMEX and the Rail Link.

The MPE site, including the Project site, is located approximately 27 km south-west of the Sydney Central Business District (CBD) and approximately 26 km west of Port Botany and includes the former Defence National Storage and Distribution Centre (DNSDC) site. The MPE site is situated within the Liverpool Local Government Area (LGA), in Sydney’s Southwest subregion, approximately 2.5 km from the Liverpool City Centre.

The MPE Project involves the development of an intermodal facility including warehouse and distribution facilities, freight village (ancillary site and operational services), stormwater infrastructure, landscaping, servicing and associated works on the eastern side of Moorebank Avenue. Stage 2 of the MPE Project involves the construction and operation of warehousing and distribution facilities on the MPE site and upgrades to approximately 2.1 km of Moorebank Avenue. Warehouses 1, 3, 4, 5, 7a and 7b are now operational. The next warehouse to be constructed is warehouse 2, which is currently planned for late 2027. Water during construction will continue to be managed in accordance with the currently approved CEMP.

BIO-ANALYSIS Pty Ltd has been commissioned by Arcadis on behalf of Tactical Group to assess stream health and water quality along Anzac Creek (the Study Area) in spring 2025.

Biodiversity Monitoring – Anzac Creek (Spring 2025)

 BIO-ANALYSIS Pty Ltd: Marine & Freshwater Ecology

Monitoring was done in accordance with a Biodiversity Monitoring Strategy (BMS) developed by Biosis (2018) to satisfy the Minister’s Conditions of Consent (CoC) B106. The BMS also includes the Stormwater Monitoring Strategy required by CoC B43 and B44.

The primary aim of monitoring was to determine whether any change in stream health or water quality occurred throughout the life of the MPE Stage 2 (MPES2) Project in accordance with the BMS and to ascertain whether these changes could be attributed to the Project works. Sampling commenced in autumn 2018 (Biosis, 2018).

2.0 METHODS


2.1 Study Area

Anzac Creek is a small tributary of the Georges River and lies entirely within the Liverpool Local Government Area. The catchment covers an area of approximately 10.6 km² (Figure 1).

The headwaters of Anzac Creek lie within the Commonwealth Department of Defence Lands in Moorebank. The creek is approximately 4 km long and highly urbanised: it flows past the suburb of Wattle Grove, underneath the M5 and Heathcote Road intersection, through the Moorebank Industrial Area and underneath Newbridge Road.

While predominantly ephemeral, Anzac Creek has been noted to hold permanent water in isolated pools (Arcadis, 2016). An unnamed first order tributary of Anzac Creek flows from south to north along the eastern boundary of the MPE Project area (GHD, 2016).

Surface water from the MPES2 site was expected to enter Anzac Creek as a licensed discharge between Site AQ4 and AQ8 (Figure 1). It was also considered likely that runoff from some areas of the MPES2 site would be collected by a vegetated dam situated within Commonwealth Department of Defence land (Biosis, 2018). Flow from this dam enters Anzac Creek upstream of Site AQ14 via a culvert (Figure 1).



Legend

- MIP East Precinct construction area
- MIP East Precinct operational area
- Warehouse layout
- Aquatic and water monitoring location
- Watercourse
- Operational rail link
- Railway

1:15,000 at A4
 GDA2020 MGA Zone 56
 Date issued: August 19, 2024
 Imagery: Maxar

ARCADIS

Figure 1. Project Location.

2.2 Sampling Dates

The dates and phases of the stream health monitoring programme for the MPES2 Project are outlined in Table 1.

Table 1. Date and information on aquatic ecology monitoring completed for the Project.

Project Phase	Event	Dates	Comments
Baseline	Autumn 2018	12&19 April 2018	Only one Baseline survey was able to be sampled in autumn 2018, due to the May 2018 bushfire.
Construction	Spring 2018	6&12 December 2018	
Construction	Autumn 2019	14&30 May 2019	Construction of culvert upstream of Site AQ1 largely completed on 30 May 2019. Site AQ12 was inaccessible to undertake Survey 2 due to restricted access.
Construction	Spring 2019	24 September 2019 21 November 2019	Warehouses 3 and 4 under construction. Moorebank Ave upgrade works ongoing.
Construction /Operation	Autumn 2020	25 May 2020 2 September 2020	Sampling required for the autumn 2020 survey season was unable to commence until late May 2020 due to COVID-19 related delays. The second survey was further delayed due to COVID related delays to receive parts required to repair the Electrofisher. Warehouses 3 and 4 were operational whilst Warehouse 5 was under construction. Moorebank Ave upgrade works ongoing.
Construction /Operation	Spring 2020	11&30 November 2020	Warehouses 3, 4 and 5 were operational. No further warehouses were being constructed at the time of monitoring
Construction /Operation	Autumn 2021	28 April 2021 11 June 2021	Warehouses 3, 4 and 5 are now operational and the location of Warehouses 6-8 have been left as compacted pads. Any water sheets off into the SED Basin and discharges into ANZAC Creek (via DP5 and DP7). No warehouses were being constructed at the time of monitoring.
Construction /Operation	Spring 2021	21 September 2021 8 November 2021	As above
Construction /Operation	Autumn 2022	5 & 31 May 2022	As above
Construction /Operation	Spring 2022	10 October 2022 30 November 2022	Following a redesign of MPE, only Warehouses 6 and 7 will be constructed within the area designated for Warehouses 6-8. Warehouse 8 will no longer be constructed. Warehouses 6&7 earthworks commenced on 9/06/22.
Construction /Operation	Autumn 2023	18 May & 3 July 2023	Warehouses 6&7 earthworks completed. It is expected that these warehouses will become operational in Q3 of 2023.

Table 1. (Cont'd)

Project Phase	Event	Dates	Comments
Construction /Operation	Spring 2023	20 September & 15 November 2023	Warehouse 7a is now operational. Operation of Warehouse 6 and 7b are expected to commence in Quarter 4 of 2023 and Quarter 2 2024.
Construction /Operation	Autumn 2024	8 & 28 May 2024	Operation of Warehouse 7b and 7a and 6 commenced in Quarter 4 of 2023 and Quarter 3 2024, respectively. The final warehouse to be constructed is warehouse 2, likely to occur in late 2025.
Construction /Operation	Spring 2024	24 September & 19 November 2024	The final warehouse to be constructed is warehouse 2, likely to occur in Q4 2025.
Construction /Operation	Autumn 2025	9 April 2025 & 23 June 2025	The final warehouse to be constructed is warehouse 2, likely to occur in late 2026.
Construction /Operation	Spring 2025	21 October & 1 December 2025	The final warehouse to be constructed is warehouse 2, likely to occur in late 2027.

2.3 Performance Measures and Indicators

Alteration to hydrology (increased stormwater inputs from both the stormwater network and surface flows from increases in non-permeable surfaces) and earthworks that have the potential to mobilise sediments into Anzac Creek were identified as potential impacts associated with the construction phase of the project (Biosis, 2018).

Biosis (2018) indicated that increased stormwater inputs to Anzac Creek could result in:

- Bed and bank scour as a result of increased volume and velocity of water during rainfall events;
- Alterations in vegetation structure as a result of altered hydrological regime;
- Introduction of sediments and pollutants via stormwater, with common pollutants including nitrogen, phosphorous, copper, aluminium and zinc.

Water Sensitive Urban Design (WSUD) measures such as onsite detention basins and rainwater gardens were incorporated into designs for the Project to mitigate impacts. A key outcome of this monitoring program was to determine whether these measures functioned as intended. Six monitoring sites (Sites AQ1, AQ4, AQ8, AQ12, AQ13 and AQ14) (Figure 1) are to be assessed in accordance with the BMS to satisfy the CoC B43, B44 and B106 (Table 2). The assessment types to be applied at each site are outlined in Table 2.

Should an indicator variable deteriorate below the range for its baseline value, a stream health investigation protocol is to be initiated under the BAEMPs Adaptive Management (Table 3). Baseline values are presented in Table 4, Table 5 and Table 6 (Results).

Table 2. Assessment types recommended for each monitoring site (Biosis, 2018).

Assessment Type	Assessment Protocol/ Indicator Variable	AQ1	AQ4	AQ8	AQ12	AQ13	AQ14
Visual	DPI Classification	√	√	√	√	√	√
	NSW AUSRIVAS	√	√	√	√	√	√
	HABSCORE	√	√	√	√	√	√
	Ephemeral Stream Assessment	√	√	√	√	√	√
Surface Water & Sediment Quality Monitoring	<i>In situ</i> water quality				√		
	Nutrient, dissolved metal & PFAS				√		
	Sediment & PFAS	√	√				√
Aquatic Macroinvertebrates	NSW AUSRIVAS & Signal2				√		
Fish	Assemblage structure				√		

Table 3. Indicator variables and adaptive management contingency measures.

Result	Potential Problem	Contingency measure
Increases in results of water quality parameters	Introduction or exacerbation of pollutants entering Anzac Creek.	Identify source and undertake corrective measures.
Reduction in results of biological monitoring	Subtle effects of construction and operation are influencing stream health within Anzac Creek.	Identify components causing decline. Assess feasibility of suitable corrective actions. If corrective measures can be implemented, these aspects are to be the focus of future monitoring. If corrective measures cannot be implemented, regulatory authority to be notified of change.
Increase scour of bed and banks of waterways	Reduction in bed and bank stability or loss of instream vegetation.	Identify point source/s of increased flow velocities or changes in stream hydraulics and discuss with project engineers to determine best methods for flow reduction or rectification of stream hydraulics

2.4 Field Methods

To fulfil the requirements of the BMS, monitoring is to be undertaken at 6 sites along Anzac Creek (Figure 1) four times annually during the pre-construction and construction phases of the Project, with the frequency reduced to twice annually during the operational phase of the Project. Surveys should take place during autumn and spring (Biosis, 2018). Sites are to be assessed using the methods outlined below and in accordance with Table 2.

Macroinvertebrate surveys were done in accordance with section 37 of the *Fisheries Management Act, 1994* using Scientific Collection Permit FP23/124 and in line with the Code of Practice Scientific Collection Activities for Fish, Invertebrates and Marine Vegetation (BIO-ANALYSIS, 2023). DPI Fisheries local inspectors were informed at least 48 hours prior to any proposed surveys as per the conditions outlined in the Scientific Collection Permit conditions.

2.4.1 Visual Stream Assessments

A visual assessment was undertaken at each site regardless of the availability of aquatic habitat (i.e. wet or dry). The condition of aquatic habitat at each site was assessed according to the *NSW Department of Primary Industries Policy and Guidelines for Fish Habitat Conservation and Management* (DPI NSW, 2013). The two key indices were habitat *type* and *class*.

Information on stream characteristics was recorded at each site in accordance with the New South Wales (NSW) Australian River Assessment System (AUSRIVAS) protocol (Turak et al., 2004). Characteristics recorded included a visual assessment of surrounding landforms, instream features, presence, extent and type of aquatic vegetation, stream substratum, potential areas of refuge during low flow periods, presence of fish habitat, presence of barriers to fish movement, indicators of point source and diffuse pollution.

HABSCORE assessments were also completed at each site, based on the presence and condition of pool substratum characteristics, pool variability, channel flow status, bank vegetation and stability, width of riparian zone, and epifaunal substrate/cover. The *CSIRO Ephemeral Stream Assessment* guideline was also used to provide an assessment of the geomorphic integrity of each site and to identify the processes operating within each site. Each site was photographed and the locations recorded with a hand-held GPS (satellite-based Global Positioning System).

2.4.2 Surface Water Quality & Sediment Monitoring

Where sufficient amounts of water were present, *in situ* water quality was measured using a Yeo-Kal 618 probe. Physico-chemical properties measured included electrical conductivity ($\mu\text{S}/\text{cm}$), dissolved oxygen (% saturation and mg/L), pH (pH units), temperature ($^{\circ}\text{C}$) and turbidity (NTU). Three replicate measures of each variable were collected from just below the water surface at each site.

Alkalinity was also determined in the field at Site AQ12, using a CHEMetrics' total alkalinity field kit.

As required by the BMS, water chemical and sediment sampling were undertaken for a range of nutrients, metals and hydrocarbons:

- Total Phosphorus (surface water only);
- Total Kjeldahl Nitrogen (TKN) (Total Organic Nitrogen + Total Ammonia) (surface water only);
- Total Nitrogen (TKN + (Nitrate + Nitrite) (surface water only);
- Dissolved metals (standard 19 relevant to aquatic assessment) (surface water);
- Total metals (standard 19 relevant to aquatic assessment) (sediment only);
- Total petroleum hydrocarbons, BTEX (benzene, toluene, ethylbenzene, trimethylbenzenes and three xylene isomers) hydrocarbons;
- PFAS: Poly-fluoroalkyl substances (including Perfluorohexane sulfonate PFHxS).

Samples were sent to the National Measurement Institute (NMI) laboratory (a NATA accredited laboratory) for analysis.

2.4.3 Aquatic Macroinvertebrates

Aquatic macroinvertebrates were required to be collected by the BMS at Site AQ12 (Biosis, 2018) using the NSW AUSRIVAS protocol (Turak et al., 2004). Biosis (2018) considered this large pool to provide reliable and valuable aquatic habitat. Stream edge habitats were sampled using a 250 µm dip net.

The contents of each net sample were placed into a white sorting tray and animals collected for a minimum period of 30 minutes. Thereafter, removals were done in 10-minute periods, up to a total of one hour (Turak et al., 2004). If no new taxa were found within a 10-minute period, removals ceased (Turak et al., 2004). The animals were collected and placed inside a labelled container and preserved with 70 % alcohol.

In the laboratory, taxa were identified to family level with the exception of Acarina (to order), Chironomidae (to sub-family), Nematoda (to phylum), Nemertea (to phylum), Oligochaeta (to class), Ostracoda (to subclass) and Polychaeta (to class). Some families of Anisoptera (dragonfly larvae) were identified to species, because they could potentially include threatened aquatic species.

2.4.4 Fish Community Survey

Fish sampling is done at Site AQ12 using a Smith Root LR-24 backpack electrofisher. The Electrofisher is used to stun fish in open water, around the edge of the pool, around snags and aquatic vegetation and any overhanging banks. All fish caught were identified and the length of up to 30 individuals of each species measured. Incidental observations such as evidence of disease were also noted before native fish species were returned to the water.

2.4.5 Data Analysis

Water quality measurements were used to assess health of the aquatic ecosystem by comparison with guideline values recommended by ANZECC² and ARMCANZ³ (2000) for the protection of lowland streams (i.e. systems at < 150 m altitude) in south-east Australia.

For aquatic macroinvertebrates, data were analysed using the appropriate AUSRIVAS predictive models developed for NSW. The ecological health of a waterway was assessed by comparing the macroinvertebrates collected at a site (i.e. Observed) to those predicted to occur (Expected) as if the site was in an undisturbed or 'reference' condition.

The principal outputs of the AUSRIVAS model include:

- Observed to Expected ratio (OE50): the ratio of the number of macroinvertebrate families collected at a site which had a predicted probability of occurrence of greater than 50 % (i.e. Observed) to the sum of the probabilities of all of the families predicted with greater than a 50 % chance of occurrence (i.e. Expected) (Ransom et al., 2004);
- BAND: for each model, the OE50 taxa ratios were divided into bands representing different levels of impairment. Band X represents a more diverse assemblage of macroinvertebrates than control sites; Band A was considered equivalent to reference condition; Band B represents sites below reference condition (i.e. significantly impaired); Band C represents sites well below reference condition (i.e. severely impaired); and Band D represents impoverished sites (i.e. extremely impaired) (Ransom et al., 2004).

The SIGNAL2 biotic index (Stream Invertebrate Grade Number Average level) developed by Chessman (2003) was also used to give an indication of water quality at the sites sampled. The SIGNAL score for a macroinvertebrate sample was calculated by averaging the pollution

² ANZECC – Australian and New Zealand Environment and Conservation Council

³ ARMCANZ – Agriculture and Resource Management Council of Australia and New Zealand

sensitivity grade numbers of the families present, which may range from 10 (most sensitive) to 1 (most tolerant). The SIGNAL2 scores from samples collected between autumn 2018 and spring 2025 were presented graphically to provide an indication of changes over time.

2.4.6 Quality Assurance/Quality Control (QA/QC)

Data collected in the field were checked for accuracy and completeness before leaving each site. In the office, field data and other records were incorporated into appropriate excel data sheets and checked. Spreadsheets were locked prior to analysis to prevent accidental overwrites or corruption.

In the laboratory, macroinvertebrate samples were identified by an appropriately qualified staff member. Data for each sample were entered into an excel spreadsheet and then checked.

3.0 RESULTS

For the spring 2025 monitoring event, sites were sampled on 21 October 2025 (Survey 1) and 1 December 2025 (Survey 2). Each site was approximately 100 m in length with their GPS co-ordinates listed in Appendix A.

Construction of the warehouses (with the exception of warehouse 2) are now complete. No construction discharges from MPE Stage 2 occurred via DP5 or DP 7 within the reporting period (i.e. after June 2025) (as per communication with Tactical).

3.1 Aquatic Habitat Characteristics

The section of Anzac Creek within the study area was not mapped as Key Fish Habitat (KFH) under the NSW DPI Key Fish Habitat mapping for the Sydney LGA (DPI 2007; Appendix A). Nevertheless, this section of Anzac Creek is ranked as TYPE 1 KFH according to the DPI (2013) classification scheme due to the presence of native aquatic plants and snags.

According to the waterway CLASS scheme, a permanent pool with freshwater aquatic vegetation situated at Site AQ12 is considered CLASS 2 KFH. The remaining reaches of Anzac Creek within the Study Area were considered to be CLASS 3 KFH despite the presence of aquatic vegetation, due to the ephemeral nature of any pools that were present (DPI, 2013).

Vegetation within the channel and banks of Anzac Creek has been classified as Parramatta Red Gum woodland in high condition (GHD, 2016).

Within the two months prior to the 2025 spring Survey 1 (21 October 2025) and Survey 2 (1 December 2025), a total of 97.6 mm and 79.2 mm rainfall was recorded respectively by the meteorological station situated near Holsworthy Aerodrome AWS Rainfall Station (Station ID: 66161) (Figure 2).

All warehouses are now operational (except warehouse 2, which is proposed to be constructed late 2027). No construction discharges from MPES2 occurred within the reporting period.




Figure 2. Rainfall (mm) measured at Holsworthy Aerodrome AWS Rainfall Station (66161) between 1 January and 31 December 2025.

Site AQ1

Site AQ1 was situated approximately 750 m downstream of the source of Anzac Creek (Figure 1), and approximately 100 m downstream of a culvert built across Anzac Creek as part of the MPES1 project. The culvert is composed of box culverts to a length of 15 m and supported one rail track and a maintenance access footway. Construction of the culvert was completed by CPB and handed over to the proponent, Qube Holdings Limited, in July 2019. The rail track has now been removed and will be rehabilitated as part of the Moorebank Avenue Realignment works however, the culverts will remain in place.

Water level was considerably lower at all sites sampled at the time of Spring 2025 Survey 1 compared to Survey 2 (Plates 1&2): isolated pools were present within the main channel at the time of Survey 1, whilst the channel was mostly submerged (to a depth of up to ~ 40 cm) at the time of Survey 2. *Gambusia (Gambusia holbrooki)* were observed.

The active channel zone remains stable (i.e., no signs of active erosion) due to dense cover of instream aquatic plants, including Slender Knotweed (*Pericaria decipiens*), Marsh Club-rush (*Bolboschoenus fluviatilis*) and Typha (*Typha* sp.), and the relatively intact woody riparian vegetation (Appendix 2).

Exotic species, including Alligator Weed (*Alternanthera philoxeroides*) and Ludwigia (*Ludwigia peruviana*), continue to be common at the edges of the channel. The channel bed consisted of fine sediment, the upper layers of which were anoxic: accumulations of green macroalgae were common at the time of Survey 2. The tree canopy was mostly comprised of *Melaleuca* spp. and *Eucalyptus* spp. (Plates 1&2).



Plate 1: AQ1 – View upstream (21/10/25)



Plate 2: AQ1 – View downstream (1/12/25)

Site AQ4

Site AQ4 was situated approximately 400 m downstream of Site AQ1 (Figure 1).

Riparian vegetation at Site AQ4, predominantly native trees, ferns including Soft Tree Fern (*Dicksonia antartica*) and stands of emergent macrophytes such as Jointed Twig Rush (*Baumea articulata*) and Twig Rush (*Baumea rubiginosa*), were mostly cleared to the edges of the creek channel to the road crossing (Plate 3&4) during winter 2025, associated with the Moorebank Avenue Realignment works⁴. The active channel zone, composed of fine sediments, is up to approximately 4 m wide, was relatively undisturbed (Plate 3&4).

⁴ The riparian corridor will be rehabilitated at the completion of the Moorebank Avenue Realignment works.

At the time of the Spring 2025 surveys, stands of *Baumea articulata* and *Baumea rubiginosa* continue to cover a large proportion of the stream channel and edges (Plates 3&4). Frogsmouth (*Philydrum lanuginosum*), *Typha* sp., *Juncus prismatocarpus*, *Carex fascicularis*, Soft Tree Fern, Heron Bristle Sedge (*Chorizandra cymbaria*) and a species of Bladderwort (*Utricularia* sp.), were also present. Water present at the time of the second survey appeared clear and free of sediment (Plate 3&4).



Plate 3: AQ4 – View downstream (21/10/25)



Plate 4: AQ4 – View across-stream (1/12/25)

Site AQ8

Site AQ8 was situated approximately 1 km downstream of Site AQ4 (Figure 1). At the time of Survey 1, small accumulations (approximately 0.25 -0.50 m²) of surface water were present to a depth of approximately 0.1 m deep. *Utricularia* sp. was present.

Stands of emergent macrophytes including Heron Bristle Sedge, Tall Spikerush (*Eleocharis sphacelata*) and Jointed Twig Rush occupied a large proportion of the stream channel (Plates 5&6). Other shorter plants, including Frogsmouth, Slender Knotweed and the introduced species, Umbrella Sedge (*Cyperus eragrostis*), have declined in abundance over the last two years, mostly likely due to shading by the taller species. Riparian vegetation continues to be dominated by *Casuarina* trees. Common Reed/Phragmites (*Phragmites australis*) and *Typha* were present at the downstream end of the site.

The stream channel at Site AQ8 (up to approximately 20 m wide) continues to be classified as stable, mostly due to the dense cover by emergent macrophytes in addition to a relatively intact, woody riparian zone (Appendix 2). Very little standing water was present within the study channel at the time of the spring 2025 surveys.



Plate 5: Site AQ8 – view downstream (21/10/25)



Plate 6: Site AQ8 – view upstream (1/12/25)

Site AQ12

Site AQ12 was situated approximately 750 km downstream of Site AQ8 (Figure 1). Similar to the findings of biodiversity surveys done since autumn 2018, a large pool (approximately 20 m wide) and a relatively diverse assemblage of aquatic plants, including submerged species, were present (Plates 7&8). Aquatic vegetation, including Slender Knotweed and dense stands of Typha, Phragmites and Tall Spike Rush have colonised a large proportion of the pool substratum (Plates 7&8). The submerged macrophyte, Ribbonweed (*Vallisneria* sp.), and floating-attached species, Entire Marshwort (*Nymphoides geminata*) (Plate 7), continue to be abundant in areas closer to the shore, which have not been colonised by tall, emergent species.

Also noted was the native perennial, *Utricularia* sp., and the small native fern, *Azolla*. *Egeria* (*Egeria densa*), which was collected close to the left-bank (facing downstream) of the pool in spring 2020, was not observed. Riparian vegetation included Casuarina, Eucalyptus and Melaleuca trees, Spiny-head Mat-rush/Basket Grass (*Lomdandra longifolia*) and *Persicaria decipiens* (Plates 7&8).

Extensive cover of vegetation within the riparian zone contributes stability to the edges of the pool at Site AQ12. An area of active erosion has been apparent at the downstream end of the pool since autumn 2020, associated with heavy rainfall and bank overflows, including at the time of the spring 2025 surveys.

Water level in the pool was approximately 0.3 m shallower than at the time of the autumn 2025 surveys. The pool edge substratum was composed primarily of fine sediment and decomposed plant material. Green filamentous macro-algae continue to be relatively abundant. Flow was not apparent at the downstream end of the pool at the time of both surveys. Water clarity was considered good.



Plate 7: Site AQ12 – view upstream (21/10/25)



Plate 8: Site AQ12 – view downstream (1/10/25)

Site AQ13

Site AQ13 was situated approximately 200 m downstream of Site AQ12 and approximately 150 m downstream from an overflow channel that enters the creek from Wattle Grove (Figure 1). A large proportion of the stream channel and edges had been recolonised by the aquatic weed, *Sagittaria platyphylla* (Sagittaria), since September 2024 (Plates 9&10). Slender Knotweed and Typha were also common. Water present was to a depth of approximately 0.5 m. Flow was not apparent. The stream channel appeared stable (Plate 10; Appendix 2).



Plate 9: Site AQ13 – view upstream (21/10/25)



Plate 10: Site AQ13 – view downstream (1/12/25)

Site AQ14

Site AQ14 was situated approximately 150 m downstream of Site AQ13 and immediately downstream of the culvert that links the dam within Commonwealth Department of Defence land to Anzac Creek (Figure 1). Flow was not apparent at the time of both spring 2025 surveys (Plates 11&12). Much of the instream vegetation scoured from much of the stream channel during autumn 2025 had recolonised the stream channel (Plates 11&12).

Typha, Slender Knotweed, River Clubrush and Whorled Pennywort/Shield Pennywort continue to be common (Plates 11&12). Sagittaria was common at the time of both survey's (Plates 11&12). This section of Anzac Creek remains mostly stable due to the presence of instream vegetation (Appendix 2). Water visibility was 'poor' at the time of first survey. Notably, some areas of instream vegetation were in poor condition, and a dead eel was observed (Plate 12).



Plate 11: Site AQ14 – view upstream (21/10/25)



Plate 12: Site AQ14 – view across-stream (1/12/25)

3.2 Water & Sediment Characteristics

3.2.1 Water Quality

Physico-chemical measurements were collected at Site AQ12 in accordance with the requirements of the BMS (cf Biosis, 2018) and at sampling sites where sufficient water was present to submerge a water quality instrument probe. The data were compared to the default trigger values (DTVs) recommended by ANZECC/ARMCANZ (2000) for the protection of slightly disturbed lowland river ecosystems in southeast Australia (Table 4).

Results from the 2025 spring surveys 1 and 2 indicated that:

- Water temperature ranged between 16.4 to 24.0 °C;
- pH (range = 6.3 to 8.2) was within the recommended DTV at all sites except Site AQ1 and AQ12 at the time of Survey 1 and AQ4 at the time of Survey 2;
- Conductivity (range = 226 to 694 µS/cm) was within the recommended DTVs at all the sites sampled;
- Dissolved oxygen (DO) measurements (range = 17.7 to 117.5 % saturation) were below the lower DTV at sites AQ1 and AQ12 during Survey 1 and at AQ4, AQ13 and AQ14 during Survey 2. Elevated levels of dissolved oxygen were recorded at AQ1 and AQ12 at the time of Survey 2;
- Turbidity levels were within the recommended DTV at all sites during spring 2025 (range = 9.1 to 49.2 NTU);
- Concentrations of total phosphorous (range = <0.05 mg/L) were within the recommended DTV (0.05 mg/L) at Site AQ12;
- Total nitrogen (range = 0.41 – 1.5 mg/L) was above the recommended DTV (0.5 mg/L) at Site AQ12 at the time of the second survey, but lower than the baseline value (i.e. 8.2 mg/L);
- Total Kjeldahl Nitrogen (TKN) (Total Organic Nitrogen + Ammonia) measured at AQ12 during both surveys was similar to the Total Nitrogen (TKN + (Nitrate + Nitrite) values, indicating that the source of nitrogen within the refuge pool was most likely organic (e.g. algae or decomposing plant material) rather than inorganic (e.g. fertilizer);
- A range of toxicants were also measured in the water between autumn 2018 (baseline) and spring 2025 (during construction) within the vicinity of Site AQ12 (Table 5&6) in accordance with the BMS (cf Biosis, 2018).

- Aluminium has commonly exceeded the DTV (80 µg/L) (i.e. 17 of 28 surveys), including at the time of the baseline survey (260 µg/L), and at the time of the first of the two spring 2025 survey's (Spring 2025 Survey 1: 410 µg/L; Survey 2: 36 µg/L);
- Cadmium exceeded the DTV (0.4 µg/L) at Site AQ12 in autumn 2019 (Survey 1: 0.49 µg/L; Survey 2: 0.41 µg/L) and autumn 2021 Survey 1 (3.8 µg/L), but not subsequently;
- Copper has commonly exceeded the DTV (1.8 µg/L) (i.e. 15 of 26 surveys, including the baseline survey (2 µg/L) but not at the time of the spring 2025 surveys;
- Zinc exceeded the DTV (i.e., 15 µg/L) during autumn 2021 (Survey 2: 20 µg/L) and autumn 2023 (Survey 2: 53 µg/L) (Table 5);
- BTEX compounds and total recoverable hydrocarbons were not detected (Table 6);
- PFOA (perfluoro-octanoic acid) has been occasionally detected but has always been well within the recommended DTV, including at the time of the spring 2025 surveys (Table 6);
- PFOS has commonly been detected, including during spring 2025 (Survey 1: 0.34 µg/L; Survey 2: 0.16 µg/L) but continues to be within the recommended DTV (Table 6).

Table 4. Mean (\pm SE) physico-chemical water quality and nutrient values recorded at the time of the Baseline (autumn 2018, $n = 1$) and the spring 2025 ($n = 3$) surveys and the appropriate Default Trigger Values (DTV). Values highlighted in bold type indicate where results were outside the recommended DTV.

Indicator Variable	DTV*	Baseline ^A	Survey 1 (21/10/25)					
			AQ1	AQ4	AQ8	AQ12	AQ13	AQ14
Temperature °C ($n=3$)	-	-	21.6 (0.0)	I/A	I/A	23.3 (0.0)	I/A	I/A
pH ($n=3$)	6.5-8.0	7.01	6.3 (0.0)	I/A	I/A	6.3 (0.0)	I/A	I/A
Conductivity (μ S/cm) ($n=3$)	125-2200	354	343 (0.9)	I/A	I/A	336 (0.0)	I/A	I/A
Dissolved Oxygen (%) ($n=3$)	85-110	62	49.1 (1.2)	I/A	I/A	44.3 (0.5)	I/A	I/A
Turbidity (NTU) ($n=3$)	<50	91	9.1 (1.2)	I/A	I/A	12.4 (0.5)	I/A	I/A
Alkalinity (mg/L) ($n=1$)	-	-	N/R	N/R	N/R	70	N/R	N/R
Total Phosphorous (mg/L) ($n=1$)	0.05	0.58	N/R	N/R	N/R	<0.05	N/R	N/R
Total Nitrogen (mg/L) ($n=1$)	0.5	8.2	N/R	N/R	N/R	1.5	N/R	N/R
Total Kjeldahl (mg/L) ($n=1$)	-	-	N/R	N/R	N/R	1.2	N/R	N/R
Indicator Variable	DTV*	Baseline ^A	Survey 2 (1/12/25)					
			AQ1	AQ4	AQ8	AQ12	AQ13	AQ14
Temperature °C ($n=3$)	-	-	22.0 (0.0)	21.2 (0.0)	I/A	24.0 (0.0)	16.4 (0.0)	17.6 (0.0)
pH ($n=3$)	6.5-8.0	7.01	7.7 (0.0)	8.2 (0.0)	I/A	6.5 (0.0)	6.4 (0.0)	6.8 (0.0)
Conductivity (μ S/cm) ($n=3$)	125-2200	354	314 (0.3)	694 (0.0)	I/A	226 (0.3)	284 (0.6)	367 (0.3)
Dissolved Oxygen (%) ($n=3$)	85-110	62	117.5 (35.9)	78.8 (0.8)	I/A	114.5 (0.3)	17.7 (0.3)	21.6 (0.2)
Turbidity (NTU) ($n=3$)	<50	91	47.7 (35.9)	18.5 (0.8)	I/A	13.4 (0.3)	49.2 (0.3)	36.3 (0.2)
Alkalinity (mg/L) ($n=1$)	-	-	N/R	N/R	N/R	30	N/R	N/R
Total Phosphorous (mg/L) ($n=1$)	0.05	0.58	N/R	N/R	N/R	<0.05	N/R	N/R
Total Nitrogen (mg/L) ($n=1$)	0.5	8.2	N/R	N/R	N/R	0.41	N/R	N/R
Total Kjeldahl (mg/L) ($n=1$)	-	-	N/R	N/R	N/R	0.41	N/R	N/R

*ANZECC/ARMCANZ (2000) – slightly disturbed systems

^A Baseline values for pH, conductivity, dissolved oxygen and turbidity were obtained from Site AQ12, whilst baseline data for phosphorous and total nitrogen were obtained from Site AQ11 (Biosis, 2018)

I/A: Insufficient Aquatic Habitat; N/R: Not Required; I/M: Instrument Malfunction. Samples were collected in the field and measured at the laboratory.

Table 5. Summary of dissolved metal compound results for Site AQ12 between autumn 2018 (Baseline) and spring 2025 (n = 1).

Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Autumn 2019 Site AQ12		Spring 2019 Site AQ12	
		April 2018	14/05/19	30/05/19	24/09/19	21/11/19
Aluminium pH >6.5	80	260	150	68	2730	280
Aluminium pH <6.5	-	-	-	-	-	-
Arsenic Total (µg/L)	42	<1	<1	<1	1.1	<1
Barium	-	2	55	34	21	32
Beryllium	-	<1	<1	<1	<1	<1
Boron	680	<50	20	17	14	14
Cadmium (µg/L)	0.4	<0.1	0.49	0.41	<0.1	<0.1
Chromium	6	<1	<1	<1	2.3	<1
Cobalt	-	<1	<1	<1	<1	<1
Copper (µg/L)	1.8	2	2	1.1	3	2.3
Iron	-	450	300	100	1650	900
Lead (µg/L)	5.6	<1	<1	<1	2.6	<1
Manganese	2500	3	33	6.2	60	47
Mercury (µg/L)	1.9 [^]	<0.1	<0.1	<0.1	0.12	<0.1
Molybdenum	-	<1	<1	<1	<1	<1
Nickel (µg/L)	13	<1	<1	N/R	1.7	1.1
Selenium Total	18	<10	<2	<1	<1	<1
Strontium	-	52	120	120	73	53
Vanadium	-	<10	<1	<1	3.8	1.4
Zinc (µg/L)	15	<5	6.8	N/R	13	14

*ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection)

[^] = inorganic mercury; N/R: not recorded

Table 5. (Cont'd). Summary of dissolved metal compound results for Site AQ12 (*n* = 1).

Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Autumn 2020 Site AQ12		Spring 2020 Site AQ12	
		April 2018	25/05/20	2/09/20	11/11/20	30/11/20
Aluminium pH >6.5	80	260	230	70	230	100
Aluminium pH <6.5	-	-	-	-	-	-
Arsenic Total (µg/L)	42	<1	<1	<1	<1	<1
Barium	-	2	31	19	36	39
Beryllium	-	<1	<1	<1	<1	<1
Boron	680	<50	21	<5	32	31
Cadmium (µg/L)	0.4	<0.1	<0.1	<0.1	<0.1	<0.1
Chromium	6	<1	<1	<1	<1	<1
Cobalt	-	<1	<1	<1	<1	<1
Copper (µg/L)	1.8	2	1.9	<1	2	1.3
Iron	-	450	620	270	460	280
Lead (µg/L)	5.6	<1	1.5	<1	<1	<1
Manganese	2500	3	19	8.8	6.9	12
Mercury (µg/L)	1.9 ^A	<0.1	<0.1	<0.1	<0.1	<0.1
Molybdenum	-	<1	1.3	<1	<1	1.1
Nickel (µg/L)	13	<1	1.1	<1	1.1	<1
Selenium Total	18	<10	<1	<1	<1	<1
Strontium	-	52	120	140	120	130
Vanadium	-	<10	<1	<1	<1	<1
Zinc (µg/L)	15	<5	8.5	3.6	5.7	2.9

*ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection)

^A = inorganic mercury; N/R: not recorded

Table 5. (Cont'd). Summary of dissolved metal compound results for Site AQ12 (n = 1).

Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Autumn 2021 Site AQ12		Spring 2021 Site AQ12	
		April 2018	28/04/21 ⁵	11/06/21	21/9/21	8/11/21
Aluminium pH >6.5	80	260	150	1260	62	200
Aluminium pH <6.5	-	-				
Arsenic Total (µg/L)	42	<1	<1	<1	<1	<1
Barium	-	2	29	<1	31	13
Beryllium	-	<1	<1	<1	<1	<1
Boron	680	<50	20	10	20	15
Cadmium (µg/L)	0.4	<0.1	3.8	<0.1	<0.1	<0.1
Chromium	6	<1	<1	1.5	<1	<1
Cobalt	-	<1	<1	<1	<1	<1
Copper (µg/L)	1.8	2	2.1	3.3	1.7	3.2
Iron	-	450	160	420	150	180
Lead (µg/L)	5.6	<1	<1	<1	<1	<1
Manganese	2500	3	6.9	4.7	10	2
Mercury (µg/L)	1.9 ^A	<0.1	<0.1	<0.1	<0.1	0.15
Molybdenum	-	<1	<1	<1	<1	<1
Nickel (µg/L)	13	<1	1.1	<1	<1	<1
Selenium Total	18	<10	<1	<1	<1	<1
Strontium	-	52	130	46	110	40
Vanadium	-	<10	<1	2.7	<1	1.9
Zinc (µg/L)	15	<5	9	20	8.3	12

*ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection)

^A = inorganic mercury; N/R: not recorded

⁵ NB Data reported here for autumn 2021 Survey 1 and Survey 2 differ from those reported in the autumn 2021 report. Data had been entered incorrectly in the autumn 2021 report but have since been corrected.

Table 5. (Cont'd). Summary of dissolved metal compound results for Site AQ12 (*n* = 1).

Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Autumn 2022 Site AQ12		Spring 2022 Site AQ12	
		April 2018	5/05/22	31/05/22	10/10/2022	30/11/2022
Aluminium pH >6.5	80	260		200	1400	93
Aluminium pH <6.5	-	-	70			
Arsenic Total (µg/L)	42	<1	<1	<1	<1	<1
Barium	-	2	18	19	15	28
Beryllium	-	<1	<1	<1	<1	<1
Boron	680	<50	21	18	26	29
Cadmium (µg/L)	0.4	<0.1	<0.1	0.13	<0.1	<0.1
Chromium	6	<1	<1	<1	1.1	<1
Cobalt	-	<1	<1	<1	<1	<1
Copper (µg/L)	1.8	2	1.4	1.5	2.6	<1
Iron	-	450	560	320	1500	350
Lead (µg/L)	5.6	<1	<1	<1	2.3	<1
Manganese	2500	3	99	5.9	9.1	16
Mercury (µg/L)	1.9 ^A	<0.1	<0.1	<0.1	<0.1	<0.1
Molybdenum	-	<1	<1	<1	<1	<1
Nickel (µg/L)	13	<1	<1	<1	<1	<1
Selenium Total	18	<10	<1	<1	<1	<1
Strontium	-	52	93	56	35	99
Vanadium	-	<10	<1	<1	2.2	<1
Zinc (µg/L)	15	<5	8	6.7	12	5.2

*ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection)

^A = inorganic mercury; N/R: not recorded

Table 5. (Cont'd). Summary of dissolved metal compound results for Site AQ12 (*n* = 1).

Indicator Variable (µg/L)	DTV*(µg/L)	Baseline Site AQ11	Autumn 2023 Site AQ12		Spring 2023 Site AQ12	
		April 2018	18/05/23	3/07/23	20/09/23	15/11/23
Aluminium pH >6.5	80	260	37	160	30	42
Aluminium pH <6.5	-	-				
Arsenic Total (µg/L)	42	<1	<1	<1	<1	<1
Barium	-	2	19	21	20	12
Beryllium	-	<1	<1	<1	<1	<1
Boron	680	<50	19	22	19	24
Cadmium (µg/L)	0.4	<0.1	0.25	0.27	<0.1	<0.1
Chromium	6	<1	<1	<1	<1	<1
Cobalt	-	<1	<1	<1	<1	<1
Copper (µg/L)	1.8	2	1.7	2.5	2.7	2.5
Iron	-	450	220	400	170	120
Lead (µg/L)	5.6	<1	<1	<1	<1	<1
Manganese	2500	3	20	40	120	11
Mercury (µg/L)	1.9 ^A	<0.1	<0.1	<0.1	<0.1	<0.1
Molybdenum	-	<1	<1	<1	<1	<1
Nickel (µg/L)	13	<1	<1	<1	<1	<1
Selenium Total	18	<10	<1	<1	<1	<1
Strontium	-	52	67	88	74	66
Vanadium	-	<10	<1	<1	<1	<1
Zinc (µg/L)	15	<5	13	53	11	2

*ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection)

^A = inorganic mercury; N/R: not recorded

Table 5. (Cont'd). Summary of dissolved metal compound results for Site AQ12 (*n* = 1).

Indicator Variable (µg/L)	DTV*(µg/L)	Baseline Site AQ11	Autumn 2024 Site AQ12		Spring 2024 Site AQ12	
		April 2018	8/05/24	28/05/24	24/09/24	19/11/24
Aluminium pH >6.5	80	260	37	170	360	290
Aluminium pH <6.5	-	-				
Arsenic Total (µg/L)	42	<1	<1	<1	<1	<1
Barium	-	2	23	18	32	18
Beryllium	-	<1	<1	<1	<1	<1
Boron	680	<50	38	32	32	25
Cadmium (µg/L)	0.4	<0.1	<0.1	<0.1	<0.1	<0.1
Chromium	6	<1	<1	<1	<1	<1
Cobalt	-	<1	<1	<1	<1	<1
Copper (µg/L)	1.8	2	1.4	1.1	1.4	2.6
Iron	-	450	310	420	1890	670
Lead (µg/L)	5.6	<1	<1	<1	<1	<1
Manganese	2500	3	5.2	19	95	36
Mercury (µg/L)	1.9 ^A	<0.1	<0.1	<0.1	<0.1	<0.1
Molybdenum	-	<1	<1	<1	<1	<1
Nickel (µg/L)	13	<1	<1	<1	<1	<1
Selenium Total	18	<10	<1	<1	<1	<1
Strontium	-	52	78	82	94	58
Vanadium	-	<10	<1	<1	2.1	1.2
Zinc (µg/L)	15	<5	13	10	12	10

*ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection)

^A = inorganic mercury; N/R: not recorded

Table 5. (Cont'd). Summary of dissolved metal compound results for Site AQ12 (n = 1).

Indicator Variable (µg/L)	DTV*(µg/L)	Baseline Site AQ11	Autumn 2025 Site AQ12		Spring 2025 Site AQ12	
		April 2018	9/04/25	23/6/25	21/10/25	1/12/25
Aluminium pH >6.5	80	260	370	43	410	36
Aluminium pH <6.5	-	-				
Arsenic Total (µg/L)	42	<1	<1	<1	1.4	<1
Barium	-	2	20	20	25	12
Beryllium	-	<1	<1	<1	<1	<1
Boron	680	<50	20	15	38	20
Cadmium (µg/L)	0.4	<0.1	<0.1	<0.1	<0.1	<0.1
Chromium	6	<1	<1	<1	<1	<1
Cobalt	-	<1	<1	<1	<1	<1
Copper (µg/L)	1.8	2	1.3	<1	1.6	<1
Iron	-	450	380	120	1040	560
Lead (µg/L)	5.6	<1	<1	<1	1	<1
Manganese	2500	3	7.8	4.8	52	28
Mercury (µg/L)	1.9 ^A	<0.1	<0.1	<0.1	<0.1	<0.1
Molybdenum	-	<1	<1	<1	<1	<1
Nickel (µg/L)	13	<1	<1	<1	<1	<1
Selenium Total	18	<10	<1	<1	<1	<1
Strontium	-	52	63	77	110	65
Vanadium	-	<10	1.6	<1	2.4	<1
Zinc (µg/L)	15	<5	7.5	6.9	8.8	<1

*ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection)

^A = inorganic mercury; N/R: not recorded

Table 6 Summary of BTEX and perfluorinated compound results (n = 1).

Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Spring 2018 Site AQ12		Autumn 2019 Site AQ12	
		April 2018	6/12/18	12/12/18	14/05/19	30/05/19
BTEX (µg/L)						
Benzene (µg/L)	1300	<1	<1	<1	<1	<1
Toluene (µg/L)	-	<2	<1	<1	<1	<1
Ethylbenzene (µg/L)	-	<2	<1	<1	<1	<1
Ortho-Xylene (µg/L)	470	<2	<1	<1	<1	<1
Perfluorinated Compounds (µg/L)						
PFHxS (µg/L)	-	0.02	0.02	0.12	0.039	0.039
PFOS (µg/L)	0.13	0.03	0.043	0.070	0.068	0.069
PFOA (µg/L)	220	<0.01	<0.01	0.011	0.011	0.010
Sum of PFHxS and PFOS	-	0.05	0.063	0.19	0.107	0.108
Sum of PFAS (WA DER List)^B	-	0.05	0.128 ^C	0.185 ^C	0.188 ^C	0.19 ^C
Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Spring 2019 Site AQ12		Autumn 2020 Site AQ12	
		April 2018	24/9/19	21/11/19	25/5/20	2/9/20
BTEX (µg/L)						
Benzene (µg/L)	1300	<1	<1	<1	<1	<1
Toluene (µg/L)	-	<2	<1	<1	<1	<1
Ethylbenzene (µg/L)	-	<2	<1	<1	<1	<1
Ortho-Xylene (µg/L)	470	<2	<1	<1	<1	<1
Perfluorinated Compounds (µg/L)						
PFHxS (µg/L)	-	0.02	0.091	0.025	0.044	0.068
PFOS (µg/L)	0.13	0.03	0.084	0.057	0.055	0.076
PFOA (µg/L)	220	<0.01	<0.01	0.013	<0.01	<0.01
Sum of PFHxS and PFOS	-	0.05	0.175	0.082	0.099	0.144
Sum of PFAS (WA DER List)^B	-	0.05	0.252 ^C	0.164 ^C	0.178 ^C	0.219 ^C

*BTEXN: ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection); PFAS suite: DEE (2016) – Freshwater (95% species protection – slightly to moderately disturbed ecosystems).

^B = PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFBS, PFHxS, PFOS, 6:2 FTS and 8:2 FTS.

^C For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01).

Table 6. (Cont'd).

Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Spring 2020 Site AQ12		Autumn 2021 Site AQ12	
		April 2018	11/11/20	30/11/20	28/04/21	11/06/21
Benzene (µg/L)	1300	<1	<1	<1	<1	<1
Toluene (µg/L)	-	<2	<1	<1	<1	<1
Ethylbenzene (µg/L)	-	<2	<1	<1	<1	<1
Ortho-Xylene (µg/L)	470	<2	<1	<1	<1	<1
PFHxS (µg/L)	-	0.02	0.026	0.041	0.065	0.011
PFOS (µg/L)	0.13	0.03	0.054	0.062	0.065	<0.02
PFOA (µg/L)	220	<0.01	0.005 ^C	0.014	<0.01	<0.01
Sum of PFHxS and PFOS	-	0.05	0.080	0.103	0.13	0.021 ^C
Sum of PFAS (WA DER List)^B	-	0.05	0.151 ^C	0.196 ^C	0.222 ^C	0.086 ^C
Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Spring 2021 Site AQ12		Autumn 2022 Site AQ12	
		April 2018	21/9/21	8/11/21	5/05/22	31/05/22
BTEX (µg/L)						
Benzene (µg/L)	1300	<1	<1	<1	<1	<1
Toluene (µg/L)	-	<2	<1	<1	<1	<1
Ethylbenzene (µg/L)	-	<2	<1	<1	<1	<1
Ortho-Xylene (µg/L)	470	<2	<1	<1	<1	<1
PFHxS (µg/L)	-	0.02	0.037	<0.01	0.044	0.039
PFOS (µg/L)	0.13	0.03	0.032	0.021	0.047	0.054
PFOA (µg/L)	220	<0.01	0.013	<0.01	<0.01	<0.01
Sum of PFHxS and PFOS	-	0.05	0.069	0.026 ^C	0.091	0.093
Sum of PFAS (WA DER List)^B	-	0.05	0.169 ^C	0.091 ^C	0.166	0.176

*BTEXN: ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection); PFAS suite: DEE (2016) – Freshwater (95 % species protection – slightly to moderately disturbed ecosystems).

^B = PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFBS, PFHxS, PFOS, 6:2 FTS and 8:2 FTS.

^C For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01).

Table 6. (Cont'd).

Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Spring 2022 Site AQ12		Autumn 2023 Site AQ12	
		April 2018	30/10/22	30/11/22	18/05/2023	3/07/2023
Benzene (µg/L)	1300	<1	<1	<1	<1	<1
Toluene (µg/L)	-	<2	<1	<1	<1	<1
Ethylbenzene (µg/L)	-	<2	<1	<1	<1	<1
Ortho-Xylene (µg/L)	470	<2	<1	<1	<1	<1
PFHxS (µg/L)	-	0.02	0.031	0.026	0.028	0.020
PFOS (µg/L)	0.13	0.03	0.030	0.044	0.040	0.024
PFOA (µg/L)	220	<0.01	<0.01	<0.01	<0.01	<0.01
Sum of PFHxS and PFOS	-	0.05	0.061	0.070	0.068	0.044
Sum of PFAS (WA DER List) ^B	-	0.05	0.126 ^C	0.135 ^C	0.145	0.122
Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Spring 2023 Site AQ12		Autumn 2024 Site AQ12	
		April 2018	20/09/23	15/11/23	8/5/24	28/5/24
Benzene (µg/L)	1300	<1	<1	<1	<1	<1
Toluene (µg/L)	-	<2	<1	<1	<1	<1
Ethylbenzene (µg/L)	-	<2	<1	<1	<1	<1
Ortho-Xylene (µg/L)	470	<2	<1	<1	<1	<1
PFHxS (µg/L)	-	0.02	0.029	0.028	0.12	0.076
PFOS (µg/L)	0.13	0.03	0.031	0.032	0.094	0.061
PFOA (µg/L)	220	<0.01	<0.01	<0.01	<0.01	<0.01
Sum of PFHxS and PFOS	-	0.05	0.060	0.060	0.214	0.137
Sum of PFAS (WA DER List) ^B	-	0.05	0.154 ^C	0.136 ^C	0.309	0.217

*BTEXN: ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection); PFAS suite: DEE (2016) – Freshwater (95% species protection – slightly to moderately disturbed ecosystems).

^B = PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFBS, PFHxS, PFOS, 6:2 FTS and 8:2 FTS.

^C For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01).

Table 6 (Cont'd).

Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Spring 2024 Site AQ12		Autumn 2025 Site AQ12	
		April 2018	24/09/24	19/11/24	9/04/25	23/6/25
Benzene (µg/L)	1300	<1	<1	<1	<1	<1
Toluene (µg/L)	-	<2	<1	<1	<1	<1
Ethylbenzene (µg/L)	-	<2	<1	<1	<1	<1
Ortho-Xylene (µg/L)	470	<2	<1	<1	<1	<1
PFHxS (µg/L)	-	0.02	0.17	0.033	0.14	0.12
PFOS (µg/L)	0.13	0.03	0.093	0.039	0.11	0.057
PFOA (µg/L)	220	<0.01	0.015	<0.01	0.012	<0.01
Sum of PFHxS and PFOS	-	0.05	0.263	0.072	0.250	0.177
Sum of PFAS (WA DER List)^B	-	0.05	0.373 ^C	0.153 ^C	0.369 ^C	0.293 ^C
Indicator Variable	DTV* (µg/L)	Baseline Site AQ11	Spring 2025 Site AQ12			
		April 2018	21/10/25	1/12/25		
Benzene (µg/L)	1300	<1	<1	<1		
Toluene (µg/L)	-	<2	<1	<1		
Ethylbenzene (µg/L)	-	<2	<1	<1		
Ortho-Xylene (µg/L)	470	<2	<1	<1		
PFHxS (µg/L)	-	0.02	0.14	0.035		
PFOS (µg/L)	0.13	0.03	0.092	0.042		
PFOA (µg/L)	220	<0.01	<0.01	<0.01		
Sum of PFHxS and PFOS	-	0.05	0.232	0.077		
Sum of PFAS (WA DER List)^B	-	0.05	0.34 ^C	0.16		

*BTEXN: ANZECC/ARMCANZ (2000) – slightly disturbed systems (90% species protection); PFAS suite: DEE (2016) – Freshwater (95% species protection – slightly to moderately disturbed ecosystems).

^B = PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFBS, PFHxS, PFOS, 6:2 FTS and 8:2 FTS.

^C For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01).

3.2.2 Sediment Characteristics

Sediment samples were collected at Site AQ1, AQ4, AQ14 between autumn 2018 (baseline) and spring 2025 (during construction) (Tables 7&8).

Results indicated that:

- Concentrations of lead measured at Site AQ1 (Survey 1: 100 mg/kg; Survey 2: 110 mg/kg) exceeded the guideline value (50 mg/L) on both sampling occasions within spring 2025. The majority (i.e. 21 of 23 times) of measurements of lead at AQ1 (range = 21 to 130 mg/kg) exceeded the threshold limit (50 mg/kg) detailed in the Interim Sediment Quality Guidelines (ISQG) (ANZECC/ARMCANZ 2000), including at the time of the baseline (91 mg/kg) survey (discussed further in Section 5.1);
- Nickel measured in sediments at Site AQ1 marginally exceeded the upper ANZECC/ARMCANZ (2000) guideline in spring 2022 (Survey 2: 25 mg/kg), spring 2023 (Survey 1: 26 mg/kg), autumn 2024 (Survey 1: 27 mg/kg), spring 2024 (Survey 1: 24 mg/kg) and spring 2025 (Survey 1: 26 mg/kg; Survey 2: 28 mg/kg);
- Concentrations of lead (56 mg/kg), nickel (23 mg/kg) and zinc (220 mg/kg) measured at AQ4 marginally exceeded the ANZECC/ARMCANZ (2000) guideline levels during Survey 1 in autumn 2022 (Table 7), but not during spring 2025;
- Zinc measured at AQ1 exceeded the upper guideline value during spring 2025 (Survey 1: 210 mg/Kg);
- Concentrations of mercury measured at AQ1 exceeded the recommended trigger level during the autumn 2022 (Survey 1: <0.2 mg/kg; Survey 2: 0.29 mg/kg) but not subsequently, including during spring 2025 (Table 7);
- A considerable spike in barium was detected at Site AQ14 in autumn 2019 (Survey 1: 902 mg/kg). There are no guideline criteria for barium in sediments or water (ANZECC/ARMCANZ 2000);
- PFOS has consistently been detected at the sites sampled (range = <0.002 to 0.044 mg/kg) but concentrations continued to be below the recommended guideline value for Urban Residential/Public Open Spaces (32 mg/kg) as well as National Parks/Areas with High Ecological Values (6.6 mg/L);

- PFAS (range = <0.001 to 0.0483 mg/kg) measured at each site continues to be similar to baseline values and below the recommended guideline value for Urban Residential/Public Open Spaces (29 mg/kg) and National Parks/Areas with High Ecological Values (1.0 mg/L) (Tables 7&8).

Table 7. Mean (\pm SE) sediment metal results (mg/L) for surveys done between autumn 2018 ($n = 1$) and spring 2025 ($n = 2$).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Autumn 2019			Spring 2019		
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	26,800	24,300 (700)	2,295 (365)	-	-	-
Antimony	-	-	-	-	<0.5	<0.5 (0)	<0.5 (0)	-	-	-
Arsenic	20	<5	<5	<5	4	6 (0.9)	1 (0.2)	3.90 (0.6)	2.75 (0.5)	2.65 (0.3)
Barium	-	110	60	<10	100	66 (4.5)	455 (447)	135 (15)	76.5 (7.5)	29.5 (1.5)
Beryllium	-	<1	1	<1	0.96	1.2 (0.0)	<0.5 (0)	1.20 (0.1)	1.01 (0.1)	<0.5 (0.00)
Boron	-	<50	<50	<50	2.9	0.8 (0.3)	<1 (0)	<1.0 (0.0)	<1.0 (0.0)	<1.0 (0.0)
Cadmium	1.5	<1	<1	<1	<0.5	<0.5 (0)	<0.5 (0)	0.43 ^A (0.2)	<0.5 (0.0)	<0.5 (0.0)
Chromium	80	23	21	3	21	23 (2.0)	3 (0.4)	21.0 (2.0)	13.5 (0.5)	6.3 (0.7)
Cobalt	-	8	6	<2	9	8 (1.9)	1 (0.1)	-	-	-
Copper	65	31	12	<5	28	11 (2.1)	2 (0.3)	30.0 (5.0)	6.1 (1.7)	9.0 (1.0)
Lead	50	91	44	<5	72	35 (0.0)	4 (0.2)	78.0 (32.0)	21.5 (0.5)	12.0 (1.0)
Manganese	-	45	69	16	32	80 (2.0)	7 (0.8)	85.0 (55.0)	50.0 (15.0)	32.5 (12.5)
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2 (0)	<0.2 (0)	<0.2 (0.0)	<0.2 (0.0)	<0.2 (0.0)
Molybdenum	-	-	-	-	2.2	1.0 (0.4)	<0.5 (0)	-	-	-
Nickel	21	14	9	<2	16	9 (0.0)	1 (0.0)	20.5 (0.5)	10.6 (1.4)	3.85 (0.2)
Selenium Total	-	<5	<5	<5	1	1 (0.0)	<0.5 (0)	2.65 (1.4)	1.59 (0.9)	0.63 ^A (0.4)
Strontium	-	-	-	-	23	17 (4.5)	1 (0.1)	-	-	-
Vanadium	-	48	54	10	36	60 (9.5)	9 (0.9)	-	-	-
Zinc	200	93	96	17	100	64 (4.0)	14 (1.5)	119 (61.5)	29 (17.5)	74 (17.0)

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Autumn 2020			Spring 2020		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	1.90 (0.2)	3.4 (0.4)	5.1 (3.1)	1.90 (0.4)	3.4 (1.2)	2.4 (0.3)
Barium	-	110	60	<10	83 (15)	63.5 (3.5)	41.3 (31.7)	87.0 (33.0)	69.5 (9.5)	37.5 (9.5)
Beryllium	-	<1	1	<1	0.72 (0.1)	0.98 (0.0)	0.5 (0.3)	0.71 (0.2)	0.79 (0.1)	<0.5 (0.0)
Boron	-	<50	<50	<50	0.85 (0.4)	0.5 (0.0)	0.5 (0.0)	1.95 (0.4)	1.25 (0.2)	0.75
Cadmium	1.5	<1	<1	<1	0.25 (0.0)	0.25 (0.0)	0.3 (0.0)	<0.05 (0.0)	<0.5 (0.0)	1.0 ^B (0.5)
Chromium	80	23	21	3	14.5 (0.5)	18.5 (0.5)	12.9 (8.2)	13.5 (3.5)	13.0 (0.0)	6.2 (0.3)
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	16.5 (0.5)	11.0 (2.0)	16.7 (12.3)	16.5 (6.5)	7.9 (0.2)	7.2 (1.2)
Lead	50	91	44	<5	71 (5.0)	33.5 (3.5)	23.5 (15.6)	53.5 (10.5)	26.0 (1.0)	11.5 (0.5)
Manganese	-	45	69	16	38.5 (0.5)	66.5 (10.5)	49.5 (38.5)	56.5 (16.5)	52.5 (4.5)	31.0 (3.0)
Mercury	0.15	<0.1	<0.1	<0.1	0.10 (0.0)	0.10 (0.0)	0.1 (0.0)	<0.2 (0.0)	<0.2 (0.0)	<0.2 (0.0)
Molybdenum	-	-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	10.7 (1.3)	8.65 (0.5)	5.4 (3.3)	11.5 (2.6)	6.5 (0.5)	2.8 (0.6)
Selenium Total	-	<5	<5	<5	0.70 (0.0)	0.44 (0.2)	0.6 (0.4)	0.63 ^B (0.4)	0.40 ^B (0.2)	<0.5 (0.0)
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	25 (1.0)	41 (2.0)	36.0 (21)	23 (5.0)	32 (5.5)	19.0 (1.0)
Zinc	200	93	96	17	78 (6.0)	144 (46.5)	111.0 (79)	86 (24)	58 (6.0)	45.5 (19.5)

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Autumn 2021			Spring 2021		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	3.65 (1.3)	6.10 (0.0)	4.30 (0.8)	14.55 (9.5)	3.5 (2.6)	2.85 (0.7)
Barium	-	110	60	<10	116.5(23.5)	99.5 (10.5)	68.0 (5.0)	74.5 (18.5)	48.0 (41.0)	84.5 (11.5)
Beryllium	-	<1	1	<1	1.20 (0.2)	0.87 (0.1)	0.50 ^A (0.2)	0.81 (0.2)	0.38 (0.4)	0.44 ^A (0.4)
Boron	-	<50	<50	<50	2.00 (0.9)	1.75 ^A (1.3)	1.40 ^A (0.9)	0.80 ^A (0.3)	<1 (0.0)	0.95 ^A (0.5)
Cadmium	1.5	<1	<1	<1	0.41 ^A (0.2)	<0.5 (0.0)	<0.5 (0.0)	<0.5 (0.0)	<0.5 (0.0)	<0.5 (0.0)
Chromium	80	23	21	3	24 (7.0)	24.5 (1.5)	13.0 (2.0)	17.5 (0.5)	12.7 (10.3)	12.0 (1.0)
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	23 (8.0)	13.5 (1.5)	12.8 (3.3)	13.0 (2.0)	6.55 (5.5)	12.3 (2.8)
Lead	50	91	44	<5	80 (50)	31.5 (2.5)	27.5 (7.5)	25.5 (4.5)	16.2 (12.9)	27.0 (7.0)
Manganese	-	45	69	16	28 (8)	150 (40)	46 (5)	95 (75)	57.1 (53)	27.5 (13.5)
Mercury	0.15	<0.1	<0.1	<0.1	<0.2 (0.0)	<0.2 (0.0)	<0.2 (0.0)	<0.2 (0.0)	<0.2 (0.0)	<0.2 (0.0)
Molybdenum	-	-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	17.5 (3.5)	9.75 (2.3)	5.85 (1.4)	10.5 (3.6)	4.1 (3.4)	7.3 (2.8)
Selenium Total	-	<5	<5	<5	1.20 (0.00)	0.88 (0.00)	0.41 (0.2)	0.88 (0.3)	0.44 ^A (0.4)	1.18 ^A (0.9)
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	10 (13)	56 (2.0)	31 (3.0)	34 (7.0)	32 (22.4)	26 (2.0)
Zinc	200	93	96	17	92 (68)	77 (14.0)	94.5 (35.5)	46 (22.0)	35 (28.2)	43 (16.0)

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Autumn 2022 (5/5/22)			Autumn 2022 (31/5/22)		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	4.3	10	6	2.9	3.6	4.6
Barium	-	110	60	<10	140	150	61	87	71	52
Beryllium	-	<1	1	<1	1.2	1.7	0.61	0.84	0.83	<0.5
Boron	-	<50	<50	<50	3.7	5	1.8	2	1.8	1
Cadmium	1.5	<1	<1	<1	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5
Chromium	80	23	21	3	23	49	11	17	20	9.9
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	24	32	14	19	14	13
Lead	50	91	44	<5	54	56	30	55	29	17
Manganese	-	45	69	16	28	320	66	25	110	41
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2	<0.2	0.29	<0.2	<0.2
Molybdenum	-	-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	17	23	5.1	13	8.8	4.2
Selenium Total	-	<5	<5	<5	3.4	3	1.3	1.1	0.68	0.57
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	37	99	31	35	46	33
Zinc	200	93	96	17	48	220	73	76	96	56

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2022 (10/10/22)			Spring 2022 (30/11/22)		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	1.9	3.6	9.8	6.1	4.1	2.1
Barium	-	110	60	<10	100	80	61	110	61	71
Beryllium	-	<1	1	<1	0.86	1	1.2	1.1	1.2	0.65
Boron	-	<50	<50	<50	4.4	2.6	4.2	1.7	<1	<1
Cadmium	1.5	<1	<1	<1	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5
Chromium	80	23	21	3	19	24	22	56	14	7.3
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	20	15	25	36	6.7	5.4
Lead	50	91	44	<5	79	32	44	62	23	12
Manganese	-	45	69	16	57	130	62	53	78	74
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
Molybdenum		-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	14	11	9.9	25	6.3	3.4
Selenium Total	-	<5	<5	<5	0.62	0.61	1.1	1	0.54	<0.5
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	24	48	67	35	40	21
Zinc	200	93	96	17	93	110	160	84	45	23

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Autumn 2023 (18/05/23)			Autumn 2023 (3/07/23)		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	26700	24500	20600	-	-	-
Antimony	-	-	-	-	<0.5	<0.5	<0.5	-	-	-
Arsenic	20	<5	<5	<5	2.8	3.1	4.6	2.9	5.1	4.2
Barium	-	110	60	<10	88	70	92	100	42	54
Beryllium	-	<1	1	<1	0.91	0.81	0.99	0.9	0.59	0.63
Boron	-	<50	<50	<50	4.5	2.2	3	2.6	<1	<1
Cadmium	1.5	<1	<1	<1	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5
Chromium	80	23	21	3	19	20	19	15	18	15
Cobalt	-	8	6	<2	7.4	7.7	6.5	-	-	-
Copper	65	31	12	<5	22	12	18	17	9.6	16
Lead	50	91	44	<5	120	25	36	37	19	32
Manganese	-	45	69	16	38	91	130	23	90	44
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
Molybdenum	-	-	-	-	1.8	0.86	0.66	-	-	-
Nickel	21	14	9	<2	14	9.9	8.3	12	5.5	6.7
Selenium Total	-	<5	<5	<5	1.3	0.79	1.1	1.6	0.53	0.68
Strontium	-	-	-	-	28	19	9.5	-	-	-
Vanadium	-	48	54	10	33	39	43	26	43	34
Zinc	200	93	96	17	100	97	77	48	54	72

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2023 (20/09/23)			Spring 2023 (15/11/23)		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	8	3.8	2.3	3.7	3.7	4.3
Barium	-	110	60	<10	140	48	42	150	79	78
Beryllium	-	<1	1	<1	1.5	0.63	<0.5	1.3	1.2	1.3
Boron	-	<50	<50	<50	6.4	<1	<1	3.7	4.2	1.2
Cadmium	1.5	<1	<1	<1	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5
Chromium	80	23	21	3	30	14	6.8	31	22	12
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	78	8.7	4.5	24	19	10
Lead	50	91	44	<5	94	20	13	87	28	17
Manganese	-	45	69	16	95	54	42	31	130	55
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
Molybdenum	-	-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	26	5.4	2.6	20	11	8.2
Selenium Total	-	<5	<5	<5	2.1	0.89	0.61	0.91	0.65	<0.5
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	51	33	20	46	40	31
Zinc	200	93	96	17	230	52	24	150	120	60

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Autumn 2024 (5/05/24)			Autumn 2024 (28/05/24)		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	5.2	1.3	1.5	3.7	3.6	5.3
Barium	-	110	60	<10	150	51	14	130	99	63
Beryllium	-	<1	1	<1	1.4	0.97	<0.5	1.1	1.5	0.94
Boron	-	<50	<50	<50	4.6	1.1	1.2	<1	<1	<1
Cadmium	1.5	<1	<1	<1	0.52	<0.5	<0.5	<0.5	<0.5	<0.5
Chromium	80	23	21	3	31	10	4.7	25	20	18
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	52	4.3	5.2	30	10	19
Lead	50	91	44	<5	100	16	7.1	73	32	38
Manganese	-	45	69	16	63	37	27	48	110	55
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
Molybdenum	-	-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	27	4.8	2.1	20	9.8	8.7
Selenium Total	-	<5	<5	<5	1.1	<0.5	<0.5	1.1	0.64	0.71
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	49	20	12	39	42	43
Zinc	200	93	96	17	200	35	44	130	73	110

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2024 (24/09/24)			Spring 2024 (19/11/24)		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	5	11	2.2	4.6	1.1	4.7
Barium	-	110	60	<10	150	110	23	97	31	75
Beryllium	-	<1	1	<1	1.5	4.1	<0.5	0.85	<0.5	0.73
Boron	-	<50	<50	<50	6	1.7	<1	6	<1	<1
Cadmium	1.5	<1	<1	<1	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5
Chromium	80	23	21	3	32	32	6.5	16	8.1	17
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	33	5.8	5.4	56	4.9	11
Lead	50	91	44	<5	100	38	9.8	53	18	28
Manganese	-	45	69	16	46	110	32	67	51	68
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
Molybdenum	-	-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	24	12	2.3	19	3.9	6.6
Selenium Total	-	<5	<5	<5	1.3	1.9	<0.5	0.87	<0.5	0.75
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	46	80	18	34	16	47
Zinc	200	93	96	17	150	47	29	230	38	73

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Autumn 2025 (9/04/25)			Autumn 2025 (23/06/25)		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	3.2	2.7	2	6	4.5	3.4
Barium	-	110	60	<10	140	60	12	180	96	65
Beryllium	-	<1	1	<1	1.1	1.1	<0.5	1.5	2	0.95
Boron	-	<50	<50	<50	4.1	2.6	<1	7.3	2.8	4.2
Cadmium	1.5	<1	<1	<1	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5
Chromium	80	23	21	3	22	16	5.6	32	20	23
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	26	8.1	4	43	4.6	35
Lead	50	91	44	<5	52	26	6.5	95	29	51
Manganese	-	45	69	16	33	79	23	54	57	50
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
Molybdenum	-	-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	17	7.1	1.6	23	7.2	9.8
Selenium Total	-	<5	<5	<5	1.5	0.8	<0.5	2	0.95	1.3
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	34	32	16	53	48	53
Zinc	200	93	96	17	62	56	39	130	39	240

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 7 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2025 (21/10/25)			Spring 2025 (1/12/25)		
		AQ1	AQ4	AQ1	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Aluminium	-	-	-	-	-	-	-	-	-	-
Antimony	-	-	-	-	-	-	-	-	-	-
Arsenic	20	<5	<5	<5	6.5	3.8	4.3	8.7	5.1	2.2
Barium	-	110	60	<10	150	50	42	170	120	17
Beryllium	-	<1	1	<1	1.2	1.1	<0.5	1.4	1.1	<0.5
Boron	-	<50	<50	<50	8.6	1.2	3.3	5.2	2.4	<1
Cadmium	1.5	<1	<1	<1	0.69	<0.5	<0.5	0.52	<0.5	<0.5
Chromium	80	23	21	3	29	16	14	36	23	6.3
Cobalt	-	8	6	<2	-	-	-	-	-	-
Copper	65	31	12	<5	81	2.6	22	59	23	7.5
Lead	50	91	44	<5	100	16	24	110	39	8.8
Manganese	-	45	69	16	52	60	140	73	160	93
Mercury	0.15	<0.1	<0.1	<0.1	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
Molybdenum	-	-	-	-	-	-	-	-	-	-
Nickel	21	14	9	<2	26	5.9	6.2	28	14	2.3
Selenium Total	-	<5	<5	<5	1.4	0.52	0.64	1.4	0.65	<0.5
Strontium	-	-	-	-	-	-	-	-	-	-
Vanadium	-	48	54	10	48	41	36	61	39	17
Zinc	200	93	96	17	210	22	160	150	140	68

*Interim Sediment Quality Guideline – Low (Trigger value) (ANZECC/ARMCANZ 2000)

^A For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (eg. <0.02 taken as 0.01)

NB Aluminium, Antimony, Molybdenum, Strontium and Vanadium were not tested for by the Spring 2019 surveys because they were not required by the BMS (cf Biosis, 2018)

Table 8. Mean (\pm SE) sediment results for perfluorinated compounds between autumn 2018 ($n = 1$) and spring 2025 ($n = 2$).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2018			Autumn 2019		
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1 ^C	AQ4	AQ14
Perfluorinated compound (mg/kg)										
PFHxS	-	0.0036	0.0007	<0.0002	0.0023 (0.00)	<0.001 (0.00)	<0.001 (0.00)	0.0037	<0.001 (0.00)	<0.001 (0.00)
PFOS	32	0.0444	0.0061	0.0005	0.0310 (0.01)	0.0049 (0.00)	<0.002 (0.00)	0.0220	0.0085 (0.01)	<0.002 (0.00)
PFOA	29	-	-	-	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001	<0.001 (0.00)	<0.001 (0.00)
Sum of PFHxS and PFOS	-	0.0480	0.0068	0.0005	0.0333 (0.01)	0.0055 ^B (0.00)	0.002 ^B (0.00)	0.0257	0.0090 ^B (0.01)	0.0015 ^B (0.00)
Sum of PFAS (WA DER List)^{A,B}	-	0.0483	0.0068	0.0005	0.0369 ^B (0.01)	0.0096 ^B (0.00)	0.0058 ^B (0.00)	0.0329	0.0150 ^B (0.01)	0.0075 ^B (0.00)
Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2019			Autumn 2020		
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
Perfluorinated compound (mg/kg)										
PFHxS	-	0.0036	0.0007	<0.0002	0.0016 (0.00)	<0.001 (0.00)	<0.001 (0.00)	0.0005 (0.00)	0.0005 (0.00)	0.0005 (0.00)
PFOS	32	0.0444	0.0061	0.0005	0.0075 (0.01)	0.0062 (0.00)	0.0028 (0.00)	0.0115 (0.00)	0.0015 (0.00)	0.0052 (0.00)
PFOA	29	-	-	-	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)
Sum of PFHxS and PFOS	-	0.0480	0.0068	0.0005	0.0231 (0.08)	0.0067 ^B (0.00)	0.0033 ^B (0.00)	0.0120 (0.00)	0.0020 (0.00)	0.0057 (0.00)
Sum of PFAS (WA DER List)^{A,B}	-	0.0483	0.0068	0.0005	0.0281 ^B (0.08)	0.0117 ^B (0.00)	0.0083 ^B (0.00)	0.0170 (0.00)	0.0070 (0.00)	0.0107 (0.00)

*DEE (2016) - Urban residential/public open spaces

^A = PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFBS, PFHxS, PFOS, 6:2 FTS and 8:2 FTS

^B For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01), the Sum of PFHxS and PFOS and the Sum of PFAS.

^C Only one survey was undertaken at Site AQ1 in autumn 2019.

Table 8 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2020			Autumn 2021		
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1 ^C	AQ4	AQ14
Perfluorinated compound (mg/kg)										
PFHxS	-	0.0036	0.0007	<0.0002	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 ^B (0.00)	<0.001 (0.00)	<0.001 (0.00)
PFOS	32	0.0444	0.0061	0.0005	0.0070 (0.00)	0.0022 ^B (0.00)	<0.002 (0.00)	0.016 (0.004)	0.006 (0.002)	0.004 (0.003)
PFOA	29	-	-	-	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)
Sum of PFHxS and PFOS	-	0.0480	0.0068	0.0005	0.0075 ^B (0.00)	0.0032 ^B (0.00)	0.0015 ^B (0.00)	0.0164 ^B (0.003)	0.0069 ^B (0.002)	0.0042 ^B (0.003)
Sum of PFAS (WA DER List)^{A,B}	-	0.0483	0.0068	0.0005	0.0125 ^B (0.00)	0.0082 ^B (0.00)	0.0065 ^B (0.00)	0.021 ^B (0.003)	0.0119 ^B (0.002)	0.0090 ^B (0.003)
Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2021			Autumn 2022		
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
PFHxS	-	0.0036	0.0007	<0.0002	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	0.0015 (0.0010)	<0.001 (0.00)	<0.001 (0.00)
PFOS	32	0.0444	0.0061	0.0005	0.0090 (0.00)	0.0030 ^B (0.00)	0.009 ^B (0.01)	0.0265 (0.0075)	0.0056 (0.0014)	0.0038 (0.0033)
PFOA	29	-	-	-	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)
Sum of PFHxS and PFOS	-	0.0480	0.0068	0.0005	0.0075 ^B (0.00)	0.0032 ^B (0.00)	0.0015 ^B (0.00)	0.0280 (0.01)	0.0056 (0.00)	0.0036 (0.0036)
Sum of PFAS (WA DER List)^{A,B}	-	0.0483	0.0068	0.0005	0.0168 ^B (0.01)	0.0089 ^B (0.00)	0.0148 ^B (0.01)	0.034 ^B (0.0075)	0.0111 ^B (0.0014)	0.0096 ^B (0.0031)

*DEE (2016) - Urban residential/public open spaces

^A = PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFBS, PFHxS, PFOS, 6:2 FTS and 8:2 FTS

^B For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01), the Sum of PFHxS and PFOS and the Sum of PFAS.

^C Only one survey was undertaken at Site AQ1 in autumn 2019.

Table 8 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2022			Autumn 2023		
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
PFHxS	-	0.0036	0.0007	<0.0002	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)
PFOS	32	0.0444	0.0061	0.0005	0.0134 (0.01)	0.0008 ^B (0.00)	<0.003 (0.00)	0.017 ^B (0.00)	0.002 ^B (0.00)	0.007 ^B (0.00)
PFOA	29	-	-	-	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)
Sum of PFHxS and PFOS	-	0.0480	0.0068	0.0005	0.0139 ^B (0.01)	0.0013 ^B (0.00)	0.0038 ^B (0.00)	0.018 ^B (0.01)	0.001 ^B (0.00)	0.004 ^B (0.00)
Sum of PFAS (WA DER List)^{A,B}	-	0.0483	0.0068	0.0005	0.0035 ^B (0.00)	0.0046 ^B (0.00)	0.0091 ^B (0.00)	0.023 ^B (0.00)	0.0075 ^B (0.001)	0.013 ^B (0.004)
Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2023			Autumn 2024		
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
PFHxS	-	0.0036	0.0007	<0.0002	<0.005 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.005 (0.00)	<0.001 (0.00)	<0.001 (0.00)
PFOS	32	0.0444	0.0061	0.0005	0.009 ^B (0.01)	0.0021 ^B (0.01)	0.0085 ^B (0.01)	0.023 ^B (0.01)	0.0022 ^B (0.01)	0.0031 ^B (0.01)
PFOA	29	-	-	-	<0.005 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.005 (0.00)	<0.001 (0.00)	<0.001 (0.00)
Sum of PFHxS and PFOS	-	0.0480	0.0068	0.0005	0.0198 ^B (0.00)	0.0034 ^B (0.00)	0.0098 ^B (0.00)	0.0340 ^B (0.00)	0.0030 ^B (0.00)	0.0043 ^B (0.00)
Sum of PFAS (WA DER List)^{A,B}	-	0.0483	0.0068	0.0005	0.0242 (0.01)	0.0076 ^B (0.00)	0.014 ^B (0.01)	0.0387 (0.01)	0.0077 ^B (0.00)	0.0089 ^B (0.00)

*DEE (2016) - Urban residential/public open spaces

^A = PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFBS, PFHxS, PFOS, 6:2 FTS and 8:2 FTS

^B For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01), the Sum of PFHxS and PFOS and the Sum of PFAS.

Table 8 (Cont'd).

Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2024			Autumn 2025		
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
PFHxS	-	0.0036	0.0007	<0.0002	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	0.001 (0.01)	0.001 (0.01)	<0.001 (0.00)
PFOS	32	0.0444	0.0061	0.0005	0.0193 (0.01)	0.0032 ^B (0.00)	<0.002 (0.00)	0.0107 (0.01)	0.0023 (0.01)	0.0045 (0.01)
PFOA	29	-	-	-	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)
Sum of PFHxS and PFOS	-	0.0480	0.0068	0.0005	0.0102 (0.01)	0.0048 ^B (0.00)	0.0023 ^B (0.00)	0.0183 ^B (0.00)	0.0057 ^B (0.00)	0.0052 ^B (0.00)
Sum of PFAS (WA DER List)^{A,B}	-	0.0483	0.0068	0.0005	0.0217 ^B (0.01)	0.0074 ^B (0.00)	0.0065 ^B (0.00)	0.0258 ^B (0.00)	0.0132 ^B (0.00)	0.0132 ^B (0.00)
Indicator Variable	Trigger Value*	Baseline (Autumn 2018)			Spring 2025					
		AQ1	AQ4	AQ14	AQ1	AQ4	AQ14	AQ1	AQ4	AQ14
PFHxS	-	0.0036	0.0007	<0.0002	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)			
PFOS	32	0.0444	0.0061	0.0005	0.0169 (0.00)	0.0036 (0.00)	0.0062 (0.00)			
PFOA	29	-	-	-	<0.001 (0.00)	<0.001 (0.00)	<0.001 (0.00)			
Sum of PFHxS and PFOS	-	0.0480	0.0068	0.0005	0.002 (0.00)	0.004 (0.00)	0.006 (0.00)			
Sum of PFAS (WA DER List)^{A,B}	-	0.0483	0.0068	0.0005	0.0308 (0.00)	0.0119 (0.00)	0.0145 (0.00)			

*DEE (2016) - Urban residential/public open spaces

^A = PFBA, PFPeA, PFHxA, PFHpA, PFOA, PFBS, PFHxS, PFOS, 6:2 FTS and 8:2 FTS

^B For any site, where a value has been recorded as less than the detection limit, it was assigned a value of half the detection limit in order to calculate the mean (e.g. <0.02 taken as 0.01), the Sum of PFHxS and PFOS and the Sum of PFAS.

3.3 Aquatic Macroinvertebrates

A total of 11 taxa were identified from edge habitat samples collected at Site AQ12 in Spring 2025 (Survey 1: 10 taxa; Survey 2: 5 taxa) (Table 9, Appendix 3). Four taxa, Chironominae (True flies), Leptoceridae (Caddis flies), Libellulidae (Dragonflies) and Physidae (aquatic snails) were collected on both sampling occasions (Appendix 3). The alien species of fish, *Gambusia*, were also collected in net samples (Appendix 3).

Site AQ12 obtained an OE50 score of 0.40 for Survey 1 and 0.20 for Survey 2 during Spring 2025 (Table 9, Figure 3), indicating that the macroinvertebrate assemblage at Site AQ12 was severely impaired (Band C) relative to reference sites selected by the AUSRIVAS model. The most recent OE50 scores were within the range of scores obtained since the baseline survey (Figure 3).


Similar to the findings of the previous surveys, taxa with > 0.80 probability of occurrence but not collected at the Anzac Creek site were Acarina (aquatic mites) during Survey 2, the aquatic beetle family, Hydrophilidae, mayfly family, Leptophlebiidae, and the aquatic bug family, Veliidae, at the time of Survey 1&2.

SIGNAL2 scores of 3.75 and 3.67 were obtained for Survey 1 and 2, respectively (Table 9). The absence of numerous taxa, including mayflies, contributed to the lower score (Table 9, Figure 4). In summary, SIGNAL 2 scores obtained for Site AQ12 have changed little over time and indicate that the macroinvertebrate assemblage at AQ12 has commonly been dominated by pollution-tolerant taxa since the commencement of sampling in autumn 2018 (Table 9, Figure 4).

Table 9. Total number of taxa, AUSRIVAS & SIGNAL 2 outputs for Site AQ12 ($n = 1$).

Survey	No Taxa	SIGNAL-2	OE50	Band
Autumn 2018	13	4.00	0.49	B
Spring 2018 – Survey 1	9	3.25	0.39	C
Spring 2018 – Survey 2	5	3.07	0.10	D
Autumn 2019 – Survey 1	10	2.69	0.41	C
Autumn 2019 – Survey 2	8	3.41	0.20	C
Spring 2019 – Survey 1	11	2.09	0.38	C
Spring 2019 – Survey 2	11	2.18	0.19	D
Autumn 2020 – Survey 1	19	3.00	0.68	B
Autumn 2020 – Survey 2	13	3.33	0.49	B
Spring 2020 – Survey 1	10	3.10	0.40	C
Spring 2020 – Survey 2	13	3.33	0.40	C
Autumn 2021 – Survey 1	13	3.38	0.49	B
Autumn 2021 – Survey 2	12	3.64	0.41	C
Spring 2021 – Survey 1	10	2.41	0.30	C
Spring 2021 – Survey 2	6	3.00	0.30	C
Autumn 2022 – Survey 1	13	3.86	0.49	B
Autumn 2022 – Survey 2	7	4.58	0.31	C
Spring 2022 – Survey 1	12	3.25	0.30	C
Spring 2022 – Survey 2	9	4.74	0.40	C
Autumn 2023 – Survey 1	7	0.30	0.29	C
Autumn 2023 – Survey 2	8	0.30	0.29	C
Spring 2023 – Survey 1	12	3.82	0.40	C
Spring 2023 – Survey 2	9	4.00	0.50	C
Autumn 2024 – Survey 1	11	3.19	0.41	C
Autumn 2024 – Survey 2	8	4.00	0.29	C
Spring 2024 – Survey 1	9	3.94	0.40	C
Spring 2024 – Survey 2	12	3.71	0.50	C
Autumn 2025 – Survey 1	6	2.64	0.19	C
Autumn 2025 – Survey 2	5	3.63	0.29	C
Spring 2025 – Survey 1	10	3.75	0.40	C
Spring 2025 – Survey 2	5	3.67	0.20	C

AUSRIVAS OE50 Scores



*Note that the bands displayed are relevant to autumn edge habitat, these being slightly different to spring

Figure 3. OE50 Taxa Scores and their respective Band Scores (B-D) for AUSRIVAS samples collected at Site AQ12 since autumn 2018.
 NB Note that the bands displayed are relevant to autumn edge habitat, these being slightly different to spring.

SIGNAL2 Scores




Figure 4. Mean (\pm) SIGNAL 2 results for Site AQ12 sampled in Anzac Creek since autumn 2018.

3.4 Fish

Three species of fish were collected while electro-fishing Site AQ12 in spring 2025, including Short-finned eel (*Anguilla australis*) (approximately 8-40 cm in length), Striped gudgeon (*Gobiomorphus australis*) and numerous Gambusia (*Gambusia holbrooki*) (Table 10). Gambusia were also abundant (Appendix 3) in net samples of aquatic macroinvertebrates.

A dead eel was observed at Site AQ14: a deep lesion was observed near the tail, similar to those observed as a result of Epizootic Ulcerative Syndrome (Red Spot Disease) (Plate 14).

In total, ten species of fish, including three introduced species, have been collected since sampling commenced in autumn 2018 (Table 10). All the species caught are common within NSW (McDowall, 1996; DPI 2006; Howell and Creese, 2010). No threatened species of fish listed under the *NSW Fisheries Management Act, 1994* or the *Environment Protection and Biodiversity Conservation Act, 1999* have been recorded.



**Plate 14: Dead eel observed at Site AQ14 at the time of Spring 2025 Survey 2.
The apparent lesion is most likely an epizootic ulcer.**

Table 10. Fish collected at Site AQ12 between autumn 2018 and spring 2025#.

Species	Common Name	A-18^	S-18	A-19	S-19	S-20	A-21	S-21	A-22	S-22	A-23	A-24	S-24	A-25	S-25
<i>Anguilla reinhardtii</i>	Long-finned eel	2	3	2	-	4	1	2	1	1	-	-	1	2	-
<i>Anguilla australis</i>	Short-finned eel	-	13	-	9	13	2	4	2	4	1	2	-		8
<i>Galaxias maculatus</i>	Common galaxias	-	-	-	-	-	-	-	8	-	-	1	1	-	-
<i>Gobiomorphus australis</i>	Striped gudgeon	28	8	3	2	-	-	-	2	2	3	1	3	2	5
<i>Hypseleotris compressa</i>	Empire gudgeon	13	-	-	-	-	-	-	-	-	1	-	-	4	-
<i>Hypseleotris cf galii</i>	Firetail gudgeon	-	-	-	1	1	-	-	-	-	-	-	1	-	-
<i>Tandanus tandanus</i>	Eel tailed catfish	-	-	-	-	-	-	-	-	-	1	-	-	-	-
<i>Carassius auratus</i> *	Goldfish	-	2	-	-	-	1	-	-	1	-	-	-	-	-
<i>Gambusia holbrooki</i> *	Gambusia	328	100's	10's	10's	100's	100's	100's	10's	100's	100's	-	80	100's	100's
<i>Misgurnus anguillicaudatus</i> *	Oriental weatherloach	-	-	-	1	-	-	-	2	1	1	1	-	-	-
<i>Retropinna semoni</i>	Australian smelt	-	-	--		-	-	-	-	-	-	-	1	-	-
Unidentified sp.	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-

^Biosis, 2018

*Introduced species

#Fish were unable to be sampled at Site AQ12 within the autumn 2020 survey period (due to instrument malfunction) or during autumn 2023 (due to the presence of extensive mats of green macro-algae).

3.5 Limitations

- Only one Baseline survey was able to be sampled in autumn 2018, due to the May 2018 bushfire (Biosis, 2018);
- Due to restricted access through the construction worksite, it was not possible to access Site AQ1 on 30 May 2019 to undertake the 2019 autumn survey 2. Whilst the collection of replicate samples at each site provides important measures of variability in habitat characteristics and concentrations of toxicants, the results from Survey 1 and subsequent surveys were within the range of results collected in the Baseline survey. Therefore, it is considered that the missing sample did not detract from being able to interpret the findings of the 2019 autumn sampling event, and that the intent and outcomes of the MPES2 monitoring survey were achieved;
- Water quality measurements collected during the biological sampling only provide a snapshot of quality at the time of sampling under the prevailing flow conditions;
- In the absence of external reference sites (i.e. similar sites but in systems not subject to the Project activities), it is not possible to account for changes in the variables examined that may occur naturally at a broader regional scale.

5.0 DISCUSSION

Stage 2 of the MPE Project involves the construction and operation of warehousing and distribution facilities on the MPE site and upgrades to approximately 2.1 kilometres of Moorebank Avenue. Warehouses 1, 3, 4, 5, 6, 7a and 7b are now operational. The next warehouse to be constructed is warehouse 2, which is currently anticipated to occur in Q4 2027. During construction of warehouse 2, water will be managed in accordance with the approved CEMP and water is discharged via the sediment (SED) Basins and into Anzac Creek (via DP5 and DP7).

5.1 Aquatic Habitat and Hydrology

Similar to the findings of surveys done since spring 2020, areas of standing water were present at the study sites and flow was apparent along some reaches during spring 2025. The majority of Anzac Creek continues to display stable environments, including areas adjacent to the Moorebank Avenue Realignment works (i.e. near Site AQ4), which commenced in June 2025. The riparian corridor will be rehabilitated following completion of the works.

Concentrations of lead in sediments collected at the most upstream site sampled on Anzac Creek (Site AQ1) (Survey 1: 100 mg/kg and Survey 2: 110 mg/kg) exceeded the guideline value (50 mg/kg) and the baseline value measured by the BAEMP survey (91 mg/kg) within spring 2025. To date, the majority (i.e. 21 of 23 times) of measurements of lead at AQ1 (range = 21 to 130 mg/kg) have exceeded the threshold limit detailed in the Interim Sediment Quality Guidelines (ISQG) (ANZECC/ARMCANZ 2000).

Nickel and zinc have also marginally exceeded the upper ANZECC/ARMCANZ (2000) guidelines, including at the time of the current survey. Site AQ1 is situated upstream of potential inputs from the Project and therefore no additional testing of heavy metals at Site AQ1 should be considered necessary.

Importantly for this survey, compounds measured in sediments at Site AQ4 remain similar to previous surveys and below the ANZECC/ARMCANZ (2000) guideline values despite removal of vegetation and disturbance of sediments within the riparian zone.

All toxicants, including total petroleum hydrocarbons and poly-fluoroalkyl substances (like PFAS and PFOS), that were monitored in sediments at the sites downstream of inputs from the MPES2 Project site (i.e., at Sites AQ4 and AQ14) in the spring 2025, remained within the appropriate guideline levels.

Measurements collected by the spring 2025 surveys indicate that additional degradation of water quality, including elevated levels of turbidity, had not occurred at Site AQ4 or at the downstream sites. It is likely that the large stands of riparian and instream aquatic plants in the downstream channel, particularly at Site AQ8, contribute to the removal of any sediments, nutrients and contaminants from the water column (Brix, 1994; Sainty and Jacobs, 2003; Feng et al., 2012; Greenway, 2010).

5.2 Water Quality

Water quality in the large refuge pool (i.e., Site AQ12) has commonly been characterised by reduced dissolved oxygen levels, elevated nitrogen, aluminium, and copper, including prior to commencement of the Project, reflecting historic and current activities (ALS, 2011; Biosis, 2018). Accumulations of green macroalgae and aquatic plants within the pool are likely to contribute to large fluctuations in dissolved oxygen and nutrient concentrations in the water column, as well as other compounds stored in the sediments into the water column.

Concentrations of total petroleum hydrocarbons and poly-fluoroalkyl substances measured during spring 2025 remain similar to baseline values and within the recommended Australian-derived guidelines for water. Additional degradation of water quality does not appear to have occurred since construction work began at Site AQ4.

5.3 Biological Monitoring

The macroinvertebrate assemblage supported by the refuge pool has consistently appeared to experience some degree of environmental stress. This is evident in the OE50 Taxa Scores and Bands, which have generally been indicative of an assemblage that is less diverse compared to reference sites selected by the AUSRIVAS model throughout the survey period. Low values of the SIGNAL 2 score and the number of macroinvertebrate types (only 5 taxa at the time of Spring 2025 - Survey 2) were also indicative of a site suffering from one or more forms of anthropogenic disturbance (see Chessman, 2003a&b).

Notably, low levels of dissolved oxygen and elevated concentrations of nitrogen have commonly been measured in the refuge pool, including during spring 2025, coinciding with excessive algal and aquatic plant growth.

Moreover, the introduced fish, *Gambusia* (*Gambusia holbrooki*), has also consistently been observed within the refuge pool. At the time of the spring 2025 survey, *Gambusia* were common in macroinvertebrate samples (50 individuals) and electrofishing samples (100's of individuals). Predation by *Gambusia* is listed as a Key Threatening Process by the NSW *Biodiversity Conservation Act 2016*, because of known effects on native frogs, freshwater fishes and aquatic macroinvertebrates.

Nevertheless, some pollution sensitive aquatic macroinvertebrates (including caddis fly and dragonfly larvae) and native species of fish continue to be collected, indicating that the creek provides important habitat for aquatic species. Of the species collected, all are common within NSW (McDowall, 1996; DPI 2006; Howell and Creese, 2010).

Notably, a dead eel was observed at Site AQ14 on 1 December 2025 (Survey 2) had a deep lesion near the tail, similar to those observed as a result of Epizootic Ulcerative Syndrome (EUS) (Red Spot Disease). EUS is caused by a fungus, *Aphanomyces invadans*, and causes red lesions (sores) and deep ulcers, making affected fish unmarketable and sometimes resulting in their death (DPI NSW, 2023). EUS is a notifiable disease in NSW and this sighting has been reported to NSW Department of Primary Industries (DPI).

Summary

This report reviews environmental monitoring results following construction activities within the Anzac Creek catchment. Most notably, riparian vegetation was cleared along an approximately 200 m section of the creek in winter 2025, associated with realignment of Moorebank Avenue. Water quality and sediment assessments found no additional degradation or increases in pollutants such as heavy metals, hydrocarbons, or poly-fluoroalkyl substances, with most measures remaining within guideline levels. Biological monitoring revealed some ongoing environmental stress in the macroinvertebrate community and the presence of the invasive *Gambusia* fish, but no measurable change in biological indices. The findings suggest that project works have not caused significant ecological impacts to date. Continued monitoring and targeted weed management are recommended.

6.0 CONCLUSION & RECOMMENDATIONS

Examination of the results from the spring 2025 monitoring event found no evidence of changes in the indicator variables (bed and bank stability, surface water and sediment quality, assemblages of aquatic macroinvertebrates and fish) that could be attributed to the Project works. Thus, in accordance with the Biodiversity Monitoring Strategy, no adaptive management contingency measures were triggered.

Recommendations include:

- Sampling of the stream health monitoring programme is repeated in autumn 2026;
- Land managers focus on containment and on-going suppression of Alligator Weed and *Ludwigia peruviana* within Anzac Creek and the riparian zone, particularly at Site AQ1, and the popular aquarium plant, *Egeria densa* (Egeria), commonly observed within the refuge pool (Site AQ12);
- In areas where riparian vegetation has recently been cleared from adjacent to Anzac Creek (i.e. within the vicinity of Site AQ4), rehabilitation works continue to be carried out to minimise sediments and contaminants becoming mobilized to the downstream environment.

7.0 REFERENCES

ALS (2011). *Assessment of the Sydney Intermodal Transport Hub, Moorebank*. Aquatic Ecology. Report prepared for Hyder Consulting Pty Ltd by Ecowise Australia Pty Ltd trading as ALS Water Resources Group.

Arcadis (2016). *Moorebank Precinct East – Stage 2 Proposal. Biodiversity Assessment Report prepared for SIMTA: Sydney Intermodal Terminal Alliance*. Part 4, Division 4.1, State Significant Development.

Australian and New Zealand Environment Conservation Council and Agriculture and Resource Management Council of Australia and New Zealand (2000). *National Water Quality Management Strategy: Australian and New Zealand Water Quality Guidelines for Fresh and Marine Water Quality*. Canberra, Australia.

Brix, H. (1994). Functions of macrophytes in constructed wetlands. *Water Science and Technology* 29: 71-78.

Biosis (2018). *Baseline Aquatic Ecological Monitoring Autumn 2018*. Report for Arcadis Authors: Stone, L. & Cable, A., Biosis Pty Ltd, Sydney. Project no. 26648.

Bureau of Meteorology [BOM] (2024). *New South Wales in 2025*.
http://www.bom.gov.au/jsp/ncc/cdio/weatherData/av?p_nccObsCode=136&p_display_type=dailyDataFile&p_startYear=&p_c=&p_stn_num=066161 (Accessed February 2026)

Chessman, B. (2003a). *SIGNAL 2 – A Scoring System for Macroinvertebrates ('Water Bugs') in Australian Rivers*. Monitoring River Health Initiative Technical Report No. 31. Commonwealth of Australia, Canberra.

Chessman, B.C. (2003b). New sensitivity grades for Australian river macroinvertebrates. *Marine and Freshwater Research*, 2003, 54: 95-103.

DPI NSW (2007). *Key Fish Habitat Maps Sydney LGA*. Website: https://www.dpi.nsw.gov.au/_data/assets/pdf_file/0009/634347/PortStephens.pdf (Accessed November 2020).

DPI NSW (2019). *NSW WeedWise*. NSW Department of Primary Industries. Website: <https://weeds.dpi.nsw.gov.au/Weeds/Alligator>. (Accessed February 2020)

Feng, W., Hatt, B. E., McCarthy, D. T., Fletcher, T. D., Deletic, A. (2012). Biofilters for stormwater harvesting: Understanding the treatment performance of key metals that pose a risk for water use. *Environmental Science and Technology* 46: 5100-5108.

GHD (2016). *Moorebank, NSW Environmental Management Plan*. Prepared for Department of Defence Former DNSDC.

Golder (2015). *Moorebank Precinct West (MPW): Site Contamination Summary Report - Stage 2 State Significant Development*. Prepared for Tactical Group on behalf of Sydney Intermodal Terminal Alliance.

Greenway, M. (2010). Wetlands and ponds for stormwater treatment in subtropical Australia: their effectiveness in enhancing biodiversity and improving water quality? *Journal of Contemporary Water Research & Education* 146: 22-38.

Howell, T. and Creese, B. (2010). *Freshwater Fish Communities of the Hunter, Manning, Karuah and Macquarie-Tuggerah Catchments: a 2004 Status Report*. Industry and Investment New South Wales, Cronulla, New South Wales.

JBS&G Australia Pty Ltd (2016). *Moorebank Precinct East (MPE) - Stage 2 Proposal: Contamination Summary Report*. Prepared for Tactical Group on behalf of Sydney Intermodal Terminal Alliance.

McDowall, R. M. (1996). *Freshwater Fishes of South-Eastern Australia*. 2nd.Edition. Reed Books, Chatswood, NSW.

Ransom, G., Coysh, J., Nichols, S. (2004). AUSRIVAS User Manual. Website: <http://ausrivas.canberra.edu.au/Bioassessment/Macroinvertebrates/Manuals> and Datasheets/User Manual. Date Retrieved: 27 November 2006.

Roberts, D. E., Church, A. G., Cummins, S. P. (1999). Invasion of *Egeria* into the Hawkesbury-Nepean River, Australia. *Journal of Aquatic Plant Management* 37: 31-34.

Sainty, G. R., Jacobs, S. W. L. (2003). *Waterplants in Australia: A Field Guide*. 4th Edn. Sainty & Associates Pty Ltd, Potts Point.

Sainty, G., McCorkelle, G., Julien, M. H. (1998). Control and Spread of alligator weed, *Alternanthera philoxeroides*, in Australia: lessons for other regions. *Wetlands Ecology Management* 5: 195–201.

Turak, E., Waddell, N., Johnstone, G. (2004). *New South Wales Australian River Assessment System (AUSRIVAS) Sampling and Processing Manual*. Department of Environment and Conservation, Sydney, Australia.

APPENDICES

Appendix 1 - GPS positions (UTMs) for stream monitoring sites (spring 2025).

Site Code	Easting	Northing
AQ1	308121	6240243
AQ4	308528	6240262
AQ8	309299	6240826
AQ12	309447	6241557
AQ13	309378	6241747
AQ14	309365	6241869

Datum: WGS 84, Zone 56H

Appendix 2 – Visual Assessment Scores

Appendix 2a – Ephemeral stream assessment results

	Autumn 2018		Spring 2018		Autumn 2019	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	88	Very Stable	75	Stable	80	Stable
AQ4	88	Very Stable	75	Stable	78	Stable
AQ8	91	Very Stable	93	Very Stable	93	Very Stable
	Spring 2019		Autumn 2020		Spring 2020	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	88	Very Stable	90	Very Stable	90	Very Stable
AQ4	80	Stable	88	Very Stable	89	Very Stable
AQ8	92	Very Stable	93	Very Stable	93	Very Stable
	Autumn 2021		Spring 2021		Autumn 2022	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	80	Very Stable	90	Very Stable	92	Very Stable
AQ4	89	Very Stable	89	Very Stable	90	Very Stable
AQ8	93	Very Stable	93	Very Stable	93	Very Stable
	Spring 2022		Autumn 2023		Spring 2023	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	92	Very Stable	88	Very Stable	88	Very Stable
AQ4	92	Very Stable	93	Very Stable	93	Very Stable
AQ8	94	Very Stable	94	Very Stable	94	Very Stable
	Autumn 2024		Spring 2024		Autumn 2025	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	94	Very Stable	92	Very Stable	90	Very Stable
AQ4	94	Very Stable	92	Very Stable	20	Very Active
AQ8	94	Very Stable	94	Very Stable	94	Very Stable

Appendix 2b – HABSCORE assessment results

	Autumn 2018		Spring 2018		Autumn 2019	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	27	Marginal	29	Marginal	32	Marginal
AQ4	28	Marginal	25	Marginal	25	Marginal
AQ8	41	Marginal	38	Marginal	38	Marginal
AQ12	55	Suboptimal	51	Suboptimal	53	Suboptimal
AQ13	21	Poor	23	Poor	21	Poor
AQ14	22	Poor	23	Poor	22	Poor
	Spring 2019		Autumn 2020		Spring 2020	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	30	Marginal	32	Marginal	27	Marginal
AQ4	26	Marginal	29	Marginal	28	Marginal
AQ8	41	Marginal	41	Marginal	41	Marginal
AQ12	51	Suboptimal	50	Suboptimal	53	Suboptimal
AQ13	19	Poor	21	Poor	22	Poor
AQ14	21	Poor	22	Poor	23	Poor
	Autumn 2021		Spring 2021		Autumn 2022	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	29	Marginal	31	Marginal	31	Marginal
AQ4	36	Marginal	38	Marginal	40	Marginal
AQ8	41	Marginal	41	Marginal	41	Marginal
AQ12	55	Suboptimal	55	Suboptimal	50	Suboptimal
AQ13	23	Poor	23	Poor	25	Poor
AQ14	24	Poor	24	Poor	25	Poor
	Spring 2022		Autumn 2023		Spring 2023	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	31	Marginal	32	Marginal	27	Marginal
AQ4	39	Marginal	40	Marginal	29	Marginal
AQ8	41	Marginal	41	Marginal	38	Marginal
AQ12	53	Suboptimal	53	Suboptimal	50	Suboptimal
AQ13	21	Poor	25	Poor	25	Poor
AQ14	25	Poor	25	Poor	25	Poor

Appendix 2b – HABSCORE assessment results

	Autumn 2024		Spring 2024		Autumn 2025	
Site	Score (%)	Category	Score (%)	Category	Score (%)	Category
AQ1	31	Marginal	27	Marginal	27	Marginal
AQ4	40	Marginal	40	Marginal	15	Poor
AQ8	41	Marginal	41	Marginal	41	Marginal
AQ12	50	Suboptimal	53	Suboptimal	53	Suboptimal
AQ13	30	Marginal	21	Poor	21	Poor
AQ14	32	Marginal	25	Poor	25	Poor
	Spring 2025					
Site	Score (%)	Category				
AQ1	40	Marginal				
AQ4	31	Marginal				
AQ8	41	Marginal				
AQ12	50	Suboptimal				
AQ13	38	Marginal				
AQ14	38	Marginal				

Appendix 3 - Macroinvertebrate taxa collected at Site AQ12 in spring 2025 using the NSW AUSRIVAS protocol. Mosquito fish were also collected in the net samples.

Taxa	Survey 1 (21 October 2025)	Survey 2 (1 December 2025)
Acariformes	22	0
Chironomidae - Chironominae	29	5
Chironomidae - Tanypodinae	2	0
Curculionidae	1	0
Isotomidae	1	0
Leptoceridae	5	1
Libellulidae	3	1
Lymnaeidae - <i>Austropeplea lessoni</i>	2	0
Pleidae	1	0
Physidae	1	1
Sphaeridae	0	1
Number of Taxa	10	5
Gambusia	10	40